



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



Elongin B (h): 293T Lysate: sc-117028

BACKGROUND

Individuals harboring germline mutations in the tumor suppressor gene von Hippel-Lindau (VHL) exhibit an increased susceptibility to a variety of tumors including renal carcinoma, hemangio-blastoma of the central nervous system and pheochromocytoma. The Elongin (SIII) complex has been identified as the functional target of the VHL protein. Elongin is a heterotrimer composed of a transcriptional active subunit designated Elongin A and two regulatory subunits designated Elongin B and Elongin C. VHL functions by binding to the Elongin B and C subunits, inhibiting the transcriptional efficacy of the Elongin complex.

REFERENCES

1. Garrett, K.P., et al. 1994. Molecular cloning of an essential subunit of RNA polymerase II elongation factor SIII. Proc. Natl. Acad. Sci. USA 91: 5237-5241.
2. Krumm, A., et al. 1995. Tumor suppression and transcription elongation: the dire consequences of changing partners. Science 269: 1400-1401.
3. Duan, D.R., et al. 1995. Inhibition of transcription elongation by the VHL tumor suppressor protein. Science 269: 1402-1406.
4. Aso, T., et al. 1995. Elongin (SIII): a multisubunit regulator of elongation by RNA polymerase II. Science 269: 1439-1443.
5. Gross, D.J., et al. 1996. Familial pheochromocytoma associated with a novel mutation in the von Hippel-Lindau gene. J. Clin. Endocrin. Metab. 81: 147-149.
6. Waber, P.G., et al. 1996. Frequent allelic loss at chromosome arm 3p is distinct from genetic alterations of the von Hippel-Lindau tumor suppressor gene in head and neck cancer. Oncogene 12: 365-369.
7. Brower, C.S., et al. 2002. Mammalian mediator subunit mMED8 is an Elongin BC-interacting protein that can assemble with CUL2 and Rbx1 to reconstitute a ubiquitin ligase. Proc. Natl. Acad. Sci. USA 99: 10353-10358.

CHROMOSOMAL LOCATION

Genetic locus: TCEB2 (human) mapping to 16p13.3.

PRODUCT

Elongin B (h): 293T Lysate represents a lysate of human Elongin B transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.


APPLICATIONS

Elongin B (h): 293T Lysate is suitable as a Western Blotting positive control for human reactive Elongin B antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

Elongin B (10C4): sc-53692 is recommended as a positive control antibody for Western Blot analysis of enhanced human Elongin B expression in Elongin B transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

DATA



Elongin B (10C4): sc-53692. Western blot analysis of Elongin B expression in non-transfected: sc-117752 (**A**) and human Elongin B transfected: sc-117028 (**B**) 293T whole cell lysates.

RESEARCH USE

For research use only, not for use in diagnostic procedures.