



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

A1Up (h): 293T Lysate: sc-117148

BACKGROUND

A1Up (Ataxin-1 ubiquitin-like-interacting protein), also known as UBQLN4 (ubiquilin 4), C1orf6 or UBIN, is a 601 amino acid protein that localizes to both the cytoplasm and the nucleus and is thought to associate with the endoplasmic reticulum (ER). Expressed at high levels in kidney, pancreas, heart, brain and skeletal muscle and at lower levels in liver, lung and placenta, A1Up functions as a homodimer that binds to signal sequences on proteins that are targeted to the ER. Additionally, A1Up is thought to link Ataxin-1 with ubiquitin/proteasome pathways, possibly assisting in the Ataxin-1-associated formation of multimeric protein complexes within the nucleus. A1Up contains one ubiquitin-like domain and one UBA domain and may be phosphorylated in response to DNA damage.

REFERENCES

1. Fogli, A., Giglio, S., Arrigo, G., Lo Nigro, C., Zollo, M., Viggiano, L., Rocchi, M., Archidiacono, N., Zuffardi, O. and Carozzo, R. 1999. Identification of two paralogous regions mapping to the short and long arms of human chromosome 2 comprising LIS1 pseudogenes. *Cytogenet. Cell Genet.* 86: 225-232.
2. Davidson, J.D., Riley, B., Burright, E.N., Duvick, L.A., Zoghbi, H.Y. and Orr, H.T. 2000. Identification and characterization of an Ataxin-1-interacting protein: A1Up, a ubiquitin-like nuclear protein. *Hum. Mol. Genet.* 9: 2305-2312.
3. Matsuda, M., Koide, T., Yoriuzzi, T., Hosokawa, N. and Nagata, K. 2001. Molecular cloning of a novel ubiquitin-like protein, UBIN, that binds to ER targeting signal sequences. *Biochem. Biophys. Res. Commun.* 280: 535-540.
4. Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 605440. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
5. Riley, B.E., Xu, Y., Zoghbi, H.Y. and Orr, H.T. 2004. The effects of the polyglutamine repeat protein Ataxin-1 on the UbL-UBA protein A1Up. *J. Biol. Chem.* 279: 42290-42301.
6. Matsuoka, S., Ballif, B.A., Smogorzewska, A., McDonald, E.R., Hurov, K.E., Luo, J., Bakalarski, C.E., Zhao, Z., Solimini, N., Lerenthal, Y., Shiloh, Y., Gygi, S.P. and Elledge, S.J. 2007. ATM and ATR substrate analysis reveals extensive protein networks responsive to DNA damage. *Science* 316: 1160-1166.
7. Li, X., Su, V., Kurata, W.E., Jin, C. and Lau, A.F. 2008. A novel connexin 43-interacting protein, CIP75, which belongs to the UbL-UBA protein family, regulates the turnover of connexin 43. *J. Biol. Chem.* 283: 5748-5759.

CHROMOSOMAL LOCATION

Genetic locus: UBQLN4 (human) mapping to 1q22.

PRODUCT

A1Up (h): 293T Lysate represents a lysate of human A1Up transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

APPLICATIONS

A1Up (h): 293T Lysate is suitable as a Western Blotting positive control for human reactive A1Up antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.