



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

hnRNP L (h): 293T Lysate: sc-117301

BACKGROUND

Heterogeneous nuclear ribonucleoproteins (hnRNPs) constitute a set of polypeptides that contribute to mRNA transcription and pre-mRNA processing as well as mature mRNA transport to the cytoplasm and translation. They also bind heterogeneous nuclear RNA (hnRNA), which are the transcripts produced by RNA Polymerase II. There are approximately 20 known hnRNP proteins, and their complexes are the major constituents of the spliceosome. The majority of hnRNP proteins are localized to the nucleus; however some shuttle between the nucleus and the cytoplasm. hnRNP I, also designated polypyrimidine tract-binding protein (PTB) and its homolog hnRNP L bind to the 3' end of introns to modulate alternative splicing mechanisms of pre-mRNAs in normal cells and the translation of several viruses, including hepatitis C virus (HCV). The human hnRNP I gene maps to chromosome 19p13.3 and encodes a protein that is localized in the nucleoplasm. hnRNP L, like hnRNP I, is also localized in the nucleoplasm.

REFERENCES

1. Badolato, J., et al. 1995. Identification and characterisation of a novel human RNA-binding protein. *Gene* 166: 323-337.
2. Siomi, H., et al. 1995. A nuclear localization domain in the hnRNP A1 protein. *J. Cell Biol.* 129: 551-560.
3. Perez, I., et al. 1997. Multiple RRM domains contribute to RNA binding specificity and affinity for polypyrimidine tract binding protein. *Biochemistry* 36: 11881-11890.
4. Hahn, B., et al. 1998. Heterogeneous nuclear ribonucleoprotein L interacts with the 3' border of the internal ribosomal entry site of hepatitis C virus. *J. Virol.* 72: 8782-8788.
5. Hahn, B., et al. 1998. Polypyrimidine tract-binding protein interacts with hnRNP L. *FEBS Lett.* 425: 401-406.
6. Li, H.P., et al. 1999. Polypyrimidine tract-binding protein binds to the leader RNA of mouse hepatitis virus and serves as a regulator of viral transcription. *J. Virol.* 73: 772-777.
7. Kim, J.H., et al. 2000. Protein-protein interaction among hnRNPs shuttling between nucleus and cytoplasm. *J. Mol. Biol.* 298: 395-405.
8. Melcak, I., et al. 2000. Nuclear pre-mRNA compartmentalization: trafficking of released transcripts to splicing factor reservoirs. *Mol. Biol. Cell* 11: 497-510.

CHROMOSOMAL LOCATION

Genetic locus: PTBP1 (human) mapping to 19p13.3.

PRODUCT

hnRNP L (h): 293T Lysate represents a lysate of human hnRNP L transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

APPLICATIONS

hnRNP L (h): 293T Lysate is suitable as a Western Blotting positive control for human reactive hnRNP L antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.