



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

RAR α (h): 293T Lysate: sc-117323

BACKGROUND

Retinoids (RA) are metabolites of vitamin A (retinol) that are important signaling molecules during vertebrate development and tissue differentiation. RAs activate the retinoic acid receptor (RAR) and retinoid X receptor (RXR) nuclear transcription factor families. Most retinoid forms activate RAR family members, whereas RXR family members are activated by 9-*cis*-RA only. RAR family members, which include RAR α , RAR β and RAR γ , have a high affinity for all transretinoic acids and belong to the same class of nuclear transcription factors as thyroid hormone receptors, vitamin D3 receptor and ecdysone receptor. RAR isoforms are expressed in distinct patterns throughout development and in the mature organism. The human RAR α gene maps to chromosome 17q21.2 and is implicated in the chromosomal translocation associated with acute promyelocytic leukemia (APL-M3). Specifically, the RAR α gene is fused with the promyelocytic leukemia (PML) gene, which encodes the fusion protein PML/RAR α . The PML/RAR α fusion protein inhibits PML-dependent apoptotic pathways and halts myeloid differentiation at the promyelocytic stage.

REFERENCES

1. Koelle, M.R., et al. 1991. The *Drosophila* EcR gene encodes an ecdysone receptor, a new member of the steroid receptor superfamily. *Cell* 67: 59-77.
2. Bhat, M.K., et al. 1994. Phosphorylation enhances the target gene sequence-dependent dimerization of thyroid hormone receptor with retinoid X receptor. *Proc. Natl. Acad. Sci. USA* 91: 7927-7931.
3. Mangelsdorf, D.J., et al. 1994. THE RETINOIDS: Biology, Chemistry, and Medicine, 2nd Edition. Sporn, M.B., Roberts, A.B. and Goodman, D.S., eds. New York: Raven Press, Ltd., 314-349.
4. Lotan, R. 1997. Roles of retinoids and their nuclear receptors in the development and prevention of upper aerodigestive tract cancers. 1997. *Environ. Health Perspect.* 4: 985-988.
5. Mark, M., et al. 1997. Genetic control of the development by retinoic acid. *C. R. Seances Soc. Biol. Fil.* 191: 77-90.
6. Randolph, T.R. 2000. Acute promyelocytic leukemia (AML-M3)—part 2: molecular defect, DNA diagnosis, and proposed models of leukemogenesis and differentiation therapy. *Clin. Lab. Sci.* 13: 106-116.
7. Wolf, G., et al. 2000. Retinoic acid induces the degradation of the leukemogenic protein encoded by the promyelocytic leukemia gene fused to the retinoic acid receptor α gene. *Nutr. Rev.* 58: 211-214.
8. Slack, J.L., et al. 2000. Current issues in the management of acute promyelocytic leukemia. *Ann. Hematol.* 79: 227-238.
9. Zhang, J.W., et al. 2000. Mechanisms of all-*trans* retinoic acid induced differentiation of acute promyelocytic leukemia cells. *J. Biosci.* 25: 275-284.

CHROMOSOMAL LOCATION

Genetic locus: RARA (human) mapping to 17q21.2.

PRODUCT

RAR α (h): 293T Lysate represents a lysate of human RAR α transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

APPLICATIONS

RAR α (h): 293T Lysate is suitable as a Western Blotting positive control for human reactive RAR α antibodies. Recommended use: 10-20 μ l per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.