



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



Hip (h2): 293T Lysate: sc-117338

BACKGROUND

The HSP 70 family is comprised of four highly conserved proteins: HSP 70, HSC 70, GRP 75 and GRP 78. These proteins serve a variety functions as molecular chaperones and aide in the assembly of multi-protein complexes. In addition to these specialized functions, the HSP 70 family may play a more general role in stabilizing protein conformation and preventing protein aggregation. HSC 70 in the mitochondrial and endoplasmic reticulum acts as ATP-driven, force generating motors that translocate proteins across organelle membranes. An HSC 70-interacting protein designated Hip, has been identified as a cochaperone in the HSC 70/HSP 40 reaction cycle. One Hip oligomer binds the ATPase domains of at least two HSC 70 molecules. This association is dependent on the activation of the HSC 70 ATPase by HSP 40. Hip functions to stabilize HSC 70 in the ADP-bound state which has high affinity for substrate protein. Through its own chaperoning activity, Hip may contribute to the substrate specificity of the HSC 70 complex.

REFERENCES

1. Martin, J., Horwich, A.L. and Hartl, F.U. 1992. Prevention of protein denaturation under heat stress by the chaperonin HSP 60. *Science* 258: 995-998.
2. Hatayama, T., Tsujioka, K., Wakatsuki, T., Kitamura, T. and Imahara, H. Effects of low culture temperature on the induction of HSP 70 mRNA and the accumulation of HSP 70 and HSP 105 in mouse FM3A cells. *J. Biochem.* 111: 484-490.
3. Bhattacharyya, T., Karnezis, A.N., Murphy, S.P., Hoang, T., Freeman, B.C., Phillips, B. and Morimoto, R.I. 1995. Cloning and subcellular localization of human mitochondrial HSP 70. *J. Biol. Chem.* 270: 1705-1710.
4. Haas, I.G. 1995. Protein-mediated protein maturation in eukaryotes. *FEBS Lett.* 369: 72-75.
5. Glick, B.S. 1995. Can HSP 70 proteins act as force-generating motors? *Cell* 80: 11-14.
6. Csermely, P., Miyata, Y., Schnaider, T. and Yahara, I. 1995. Autophosphorylation of GRP 94 (endoplasmin). *J. Biol. Chem.* 270: 6381-6388.
7. Höhfeld, J., Minami, Y. and Hartl, F.U. 1995. Hip, a novel cochaperone involved in the eukaryotic HSC 70/ HSP 40 reaction cycle. *Cell* 83: 589-598.

CHROMOSOMAL LOCATION

Genetic locus: ST13 (human) mapping to 22q13.2.

PRODUCT

Hip (h2): 293T Lysate represents a lysate of human Hip transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

APPLICATIONS

Hip (h2): 293T Lysate is suitable as a Western Blotting positive control for human reactive Hip antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.