



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



Myo-inositol oxygenase (h2): 293T Lysate: sc-117367

BACKGROUND

Myo-inositol oxygenase (MIOX), also known as ALDRL6, is a renal-specific member of the aldo-keto reductase family. It catalyzes the first committed step in the Myo-inositol metabolism pathway and is widely distributed in mammalian tissues. Human Myo-inositol oxygenase shares 91% and 96% sequence homology with mouse and pig Myo-inositol oxygenase homologs, respectively. Myo-inositol oxygenase is responsible for the oxidative cleavage of Myo-inositol (MI) and its epimer D-chiro inositol (DCI) to D-glucuronate. The dioxygen-dependent cleavage of the C1-C6 bond in Myo-inositol is accomplished through the utilization of the Fe^{II}/Fe^{III} binuclear iron center of MIOX. Myo-inositol oxygenase has also been implicated in complications of diabetes, including diabetic nephropathy.

REFERENCES

1. Yang, Q., Dixit, B., Wada, J., Tian, Y., Wallner, E.I., Srivastva, S.K. and Kanwar, Y.S. 2000. Identification of a renal-specific oxido-reductase in newborn diabetic mice. *Proc. Natl. Acad. Sci. USA* 97: 9896-9901.
2. Arner, R.J., Prabhu, K.S., Thompson, J.T., Hildenbrandt, G.R., Liken, A.D. and Reddy, C.C. 2001. Myo-inositol oxygenase: molecular cloning and expression of a unique enzyme that oxidizes Myo-inositol and D-chiro-inositol. *Biochem. J.* 360: 313-320.
3. Lorence, A., Chevone, B.I., Mendes, P. and Nessler, C.L. 2004. Myo-inositol oxygenase offers a possible entry point into plant ascorbate biosynthesis. *Plant Physiol.* 134: 1200-1205.
4. Arner, R.J., Prabhu, K.S., Krishnan, V., Johnson, M.C. and Reddy, C.C. 2005. Expression of Myo-inositol oxygenase in tissues susceptible to diabetic complications. *Biochem. Biophys. Res. Commun.* 339: 816-820.
5. Prabhu, K.S., Arner, R.J., Vunta, H. and Reddy, C.C. 2005. Up-regulation of human Myo-inositol oxygenase by hyperosmotic stress in renal proximal tubular epithelial cells. *J. Biol. Chem.* 280: 19895-19901.
6. Brown, P.M., Caradoc-Davies, T.T., Dickson, J.M., Cooper, G.J., Loomes, K.M. and Baker, E.N. 2006. Purification, crystallization and preliminary crystallographic analysis of mouse Myo-inositol oxygenase. *Acta Crystallogr. Sect. F Struct. Biol. Cryst. Commun.* 62: 811-813.
7. Xing, G., Hoffart, L.M., Diao, Y., Prabhu, K.S., Arner, R.J., Reddy, C.C., Krebs, C. and Bollinger, J.M. 2006. A coupled dinuclear iron cluster that is perturbed by Myo-inositol oxygenase. *Biochemistry* 45: 5393-5401.
8. Xing, G., Barr, E.W., Diao, Y., Hoffart, L.M., Prabhu, K.S., Arner, R.J., Reddy, C.C., Krebs, C. and Bollinger, J.M. 2006. Oxygen activation by a mixed-valent, diiron(II/III) cluster in the glycol cleavage reaction catalyzed by Myo-inositol oxygenase. *Biochemistry* 45: 5402-5412.
9. Xing, G., Diao, Y., Hoffart, L.M., Barr, E.W., Prabhu, K.S., Arner, R.J., Reddy, C.C., Krebs, C. and Bollinger, J.M. 2006. Evidence for C-H cleavage by an iron-superoxide complex in the glycol cleavage reaction catalyzed by Myo-inositol oxygenase. *Proc. Natl. Acad. Sci. USA* 103: 6130-6135.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

CHROMOSOMAL LOCATION

Genetic locus: MIOX (human) mapping to 22q13.33.

PRODUCT

Myo-inositol oxygenase (h2): 293T Lysate represents a lysate of human Myo-inositol oxygenase transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

APPLICATIONS

Myo-inositol oxygenase (h2): 293T Lysate is suitable as a Western Blotting positive control for human reactive Myo-inositol oxygenase antibodies. Recommended use: 10-20 µl per lane.


Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

Myo-inositol oxygenase (D-11): sc-271512 is recommended as a positive control antibody for Western Blot analysis of enhanced human Myo-inositol oxygenase expression in Myo-inositol oxygenase transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:
 1) Western Blotting: use m-IgG_κ BP-HRP: sc-516102 or m-IgG_κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

DATA



Myo-inositol oxygenase (D-11): sc-271512. Western blot analysis of Myo-inositol oxygenase expression in non-transfected: sc-117752 (**A**) and human Myo-inositol oxygenase transfected: sc-117367 (**B**) 293T whole cell lysates.

Myo-inositol oxygenase (E-11): sc-376080. Western blot analysis of Myo-inositol oxygenase expression in non-transfected: sc-117752 (**A**) and human Myo-inositol oxygenase transfected: sc-117367 (**B**) 293T whole cell lysates.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.