



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# 4E-BP1 (m): 293T Lysate: sc-118030

## BACKGROUND

The translation of proteins from eukaryotic mRNA is initiated by the multi-subunit complex eIF-4F, which associates with the mRNA 5' cap structure. eIF-4E, a component of eIF-4F, is responsible for binding to the 5' cap structure and for the assembly of the eIF-4F complex. The regulatory protein 4E-BP1, also referred to as PHAS-I, inhibits eIF-4E function. Phosphorylation of 4E-BP1 by S6 kinase p70, MAP kinases or PKCs causes the disassociation of 4E-BP1 from eIF-4E, promoting translation. A protein that is functionally related to 4E-BP1, designated 4E-BP2, also associates with eIF-4E.

## REFERENCES

1. Lin, T.A., et al. 1994. PHAS-I as a link between mitogen-activated protein kinase and translation initiation. *Science* 266: 653-656.
2. Rau, M., et al. 1996. A reevaluation of the cap-binding protein, eIF4E, as a rate-limiting factor for initiation of translation in reticulocyte lysate. *J. Biol. Chem.* 271: 8983-8990.
3. Whalen, S.G., et al. 1996. Phosphorylation of eIF-4E on Serine 209 by protein kinase C is inhibited by the translational repressors, 4E-binding proteins. *J. Biol. Chem.* 271: 11831-11837.
4. Diggle, T.A., et al. 1996. Both Rapamycin-sensitive and -insensitive pathways are involved in the phosphorylation of the initiation factor-4E-binding protein (4E-BP1) in response to Insulin in rat epididymal fat-cells. *Biochem. J.* 316: 447-453.
5. Beretta, L., et al. 1996. Rapamycin blocks the phosphorylation of 4E-BP1 and inhibits cap-dependent initiation of translation. *EMBO J.* 15: 658-64.
6. Mendez, R., et al. 1996. Stimulation of protein synthesis, eukaryotic translation initiation factor 4E phosphorylation, and PHAS-I phosphorylation by Insulin requires Insulin receptor substrate 1 and phosphatidylinositol 3-kinase. *Mol. Cell. Biol.* 16: 2857-2864.

## CHROMOSOMAL LOCATION

Genetic locus: Eif4ebp1 (mouse) mapping to 8 A2.

## PRODUCT

4E-BP1 (m): 293T Lysate represents a lysate of mouse 4E-BP1 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

## APPLICATIONS

4E-BP1 (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive 4E-BP1 antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

4E-BP1 (P-1): sc-9977 is recommended as a positive control antibody for Western Blot analysis of enhanced mouse 4E-BP1 expression in 4E-BP1 transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).


## RESEARCH USE

For research use only, not for use in diagnostic procedures.


## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-mouse IgG-HRP: sc-2005 (dilution range: 1:2000-1:32,000) or Cruz Marker™ compatible goat anti-mouse IgG-HRP: sc-2031 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048.

## DATA



4E-BP1 (P-1): sc-9977. Western blot analysis of 4E-BP1 expression in non-transfected: sc-117752 (A) and mouse 4E-BP1 transfected: sc-118030 (B) 293T whole cell lysates.



4E-BP1 (C-19): sc-6024. Western blot analysis of 4E-BP1 expression in non-transfected: sc-117752 (A) and mouse 4E-BP1 transfected: sc-118030 (B) 293T whole cell lysates.

## STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.