



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

ACAT-2 (m2): 293T Lysate: sc-118190

BACKGROUND

ACAT-1 (acetyl-coenzyme A acetyltransferase 1) is a mitochondrial enzyme involved in the formation and degradation of ketone bodies and is necessary for the proper metabolic processing of isoleucine. Rare defects in the gene encoding ACAT-1 lead to β -ketothiolase deficiency, which is characterized by ketoacidotic attacks. ACAT-2 (acetyl-coenzyme A acetyltransferase 2) is considered a cytosolic protein and is crucial for cholesterol synthesis. Specifically, both acetoacetyl-CoA specific thiolases, ACAT-1 and ACAT-2 catalyze the formation of acetoacetyl-CoA from two acetyl-CoA molecules. These enzymes are also capable of the reverse reaction, the cleavage of acetoacetyl-CoA into two acetyl-CoA molecules.

REFERENCES

1. Groot, C.J., Haan, G.L., Hulstaert, C.E. and Hoopes, F.A. 1977. A patient with severe neurologic symptoms and acetoacetyl-CoA thiolase deficiency. *Pediatr. Res.* 11: 1112-1116.
2. Willison, K., Kelly, A., Dudley, K., Goodfellow, P., Spurr, N., Groves, V., Gorman, P., Sheer, D. and Trowsdale, J. 1987. The human homologue of the mouse t-complex gene, TCP1, is located on chromosome 6 but is not near the HLA region. *EMBO J.* 6: 1967-1974.
3. Igual, J.C., González-Bosch, C., Dopazo, J. and Pérez-Ortín, J.E. 1992. Phylogenetic analysis of the thiolase family. Implications for the evolutionary origin of peroxisomes. *J. Mol. Evol.* 35: 147-155.
4. Ashworth, A. 1993. Two acetyl-CoA acetyltransferase genes located in the t-complex region of mouse chromosome 17 partially overlap the *Tcp-1* and *Tcp-1x* genes. *Genomics* 18: 195-198.
5. Song, X.Q., Fukao, T., Yamaguchi, S., Miyazawa, S., Hashimoto, T. and Orii, T. 1994. Molecular cloning and nucleotide sequence of complementary DNA for human hepatic cytosolic acetoacetyl-coenzyme A thiolase. *Biochem. Biophys. Res. Commun.* 201: 478-485.
6. Masuno, M., Fukao, T., Song, X.Q., Yamaguchi, S., Orii, T., Kondo, N., Imaizumi, K. and Kuroki, Y. 1996. Assignment of the human cytosolic acetoacetyl-coenzyme A thiolase (ACAT2) gene to chromosome 6q25.3-q26. *Genomics* 36: 217-218.
7. Haapalainen, A.M., Meriläinen, G., Pirilä, P.L., Kondo, N., Fukao, T. and Wierenga, R.K. 2007. Crystallographic and kinetic studies of human mitochondrial acetoacetyl-CoA thiolase: the importance of potassium and chloride ions for its structure and function. *Biochemistry* 46: 4305-4321.

CHROMOSOMAL LOCATION

Genetic locus: *Acat2* (mouse) mapping to 17 A1.

PRODUCT

ACAT-2 (m2): 293T Lysate represents a lysate of mouse ACAT-2 transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

APPLICATIONS

ACAT-2 (m2): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive ACAT-2 antibodies. Recommended use: 10-20 μ l per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.