



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

AdoMetDC (m2): 293T Lysate: sc-118260

BACKGROUND

Polyamines are compounds that have two or more primary amino groups and are important to cellular processes, such as cellular growth, proliferation and tumor promotion. AdoMetDC (adenosylmethionine decarboxylase 1), also known as S-adenosylmethionine decarboxylase proenzyme (SAMDC) or AMD1, is a 334 amino acid protein which is an important intermediate enzyme in polyamine biosynthesis pathways. Using a pyruvoyl group as a cofactor, AdoMetDC catalyzes the conversion of S-adenosyl-L-methionine to (5-deoxy-5-adenosyl) (3-aminopropyl)-methylsulfonium salt and carbon dioxide. AdoMetDC is synthesized as an inactive proenzyme that undergoes self-maturation to form two non-identical subunits designated α and β . Active AdoMetDC forms a heterotetramer of two α chains and two β chains. Both AdoMetDC proenzyme processing and mature AdoMetDC catalytic activity are stimulated by putrescine, while catalytic activity is inhibited by iodoacetic acid.

REFERENCES

- Ekstrom, J.L., et al. 2001. Structure of a human S-adenosylmethionine decarboxylase self-processing ester intermediate and mechanism of putrescine stimulation of processing as revealed by the H243A mutant. *Biochemistry* 40: 9495-9504.
- Tolbert, W.D., et al. 2003. Mechanism of human S-adenosylmethionine decarboxylase proenzyme processing as revealed by the structure of the S68A mutant. *Biochemistry* 42: 2386-2395.
- Yerlikaya, A., et al. 2004. S-adenosylmethionine decarboxylase degradation by the 26S Proteasome is accelerated by substrate-mediated transamination. *J. Biol. Chem.* 279: 12469-12478.
- Lam, K., et al. 2005. HSG cells differentiated by culture on extracellular matrix involves induction of S-adenosylmethionine decarboxylase and ornithine decarboxylase. *J. Cell. Physiol.* 203: 353-361.
- Kim, J.S., et al. 2006. S-Adenosylmethionine decarboxylase partially regulates cell growth of HL-60 cells by controlling the intracellular ROS level: Early senescence and sensitization to γ -radiation. *Arch. Biochem. Biophys.* 456: 58-70.
- Guidotti, A., et al. 2007. S-adenosylmethionine and DNA methyltransferase-1 mRNA overexpression in psychosis. *Neuroreport* 18: 57-60.
- Bale, S., et al. 2008. Structural basis for putrescine activation of human S-adenosylmethionine decarboxylase. *Biochemistry* 47: 13404-13417.

CHROMOSOMAL LOCATION

Genetic locus: Amd1 (mouse) mapping to 10 B1.

PRODUCT

AdoMetDC (m2): 293T Lysate represents a lysate of mouse AdoMetDC transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

APPLICATIONS

AdoMetDC (m2): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive AdoMetDC antibodies. Recommended use: 10-20 μ l per lane.


Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

AdoMetDC (F-10): sc-398482 is recommended as a positive control antibody for Western Blot analysis of enhanced mouse AdoMetDC expression in AdoMetDC transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

DATA



AdoMetDC (F-10): sc-398482. Western blot analysis of AdoMetDC expression in non-transfected: sc-117752 (A) and mouse AdoMetDC transfected: sc-118260 (B) 293T whole cell lysates.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.