



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Aladin (m2): 293T Lysate: sc-118322

BACKGROUND

Aladin (Adracalin) belongs to a family of WD repeat-containing proteins. These proteins have a wide variety of functions, including signal transduction regulation, RNA processing and transcription. Aladin plays a role in peripheral and central nervous system development. It is widely expressed, with the highest expression seen in pituitary gland, corpus callosum, cerebellum, adrenal gland and gastrointestinal structures. Defects in Aladin cause the autosomal recessive disorder achalasia-addisonianism-alacrima (triple A) syndrome. Triple A syndrome is characterized by achalasia, alacrima and adrenocorticotropin-resistant adrenal insufficiency. Robust expression in neural systems associated with cognitive, motor and sensory functions is consistent with the myriad of symptoms experienced by patients with triple A syndrome.

REFERENCES

1. Tullio-Pelet, A., et al. 2000. Mutant WD-repeat protein in triple A syndrome. *Nat. Genet.* 26: 332-335.
2. Katsumata, N., et al. 2002. Analysis of the AAAS gene in a Japanese patient with triple A syndrome. *Endocr. J.* 49: 49-53.
3. Houlden, H., et al. 2002. Clinical and genetic characterization of families with triple A (Allgrove) syndrome. *Brain* 125: 2681-2690.
4. Cronshaw, J.M., et al. 2003. The nuclear pore complex protein Aladin is mislocalized in triple A syndrome. *Proc. Natl. Acad. Sci. USA* 100: 5823-5827.
5. Salehi, M., et al. 2005. The diagnosis of adrenal insufficiency in a patient with Allgrove syndrome and a novel mutation in the Aladin gene. *Metabolism* 54: 200-205.
6. Storr, H.L., et al. 2005. Identification of the sites of expression of triple A syndrome mRNA in the rat using *in situ* hybridisation. *Neuroscience* 131: 113-123.

CHROMOSOMAL LOCATION

Genetic locus: Aaas (mouse) mapping to 15 F3.

PRODUCT

Aladin (m2): 293T Lysate represents a lysate of mouse Aladin transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

Aladin (m2): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive Aladin antibodies. Recommended use: 10-20 µl per lane.


Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

Aladin (3E9): sc-100321 is recommended as a positive control antibody for Western Blot analysis of enhanced mouse Aladin expression in Aladin transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:
1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

DATA



Aladin (3E9): sc-100321. Western blot analysis of Aladin expression in non-transfected: sc-117752 (A) and mouse Aladin transfected: sc-118322 (B) 293T whole cell lysates.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.