



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

AQP4 (m): 293T Lysate: sc-118504

BACKGROUND

In skeletal muscle, AQP4 (aquaporin 4 also known as mercurial insensitive water channel), localizes to the sarcolemma of fast-twitch muscle fibers. Aquaporins (AQPs) are a large family of integral membrane water transport channel proteins that facilitate the transport of water through the cell membrane. This function is conserved in animals, plants and bacteria. Many isoforms of aquaporin have been identified in mammals, designated AQPO through AQP10. Aquaporins are widely distributed and it is not uncommon for more than one type of AQP to be present in the same cell. Although most aquaporins are only permeable to water, AQP3, AQP7, AQP9 and one of the two AQP10 transcripts are also permeable to urea and glycerol. AQP2 is the only water channel that is activated by vasopressin to enhance water reabsorption in the kidney collecting duct. Aquaporins are involved in renal water absorption, generation of pulmonary secretions, lacrimation and the secretion and reabsorption of cerebrospinal fluid and aqueous humor.

REFERENCES

1. Preston, G.M., et al. 1991. Isolation of the cDNA for erythrocyte integral membrane protein of 28 kDa: member of an ancient channel family. *Proc. Natl. Acad. Sci. USA* 88: 11110-11114.
2. Deen, P.M., et al. 1994. Requirement of human renal water channel aquaporin 2 for vasopressin-dependent concentration of urine. *Science* 264: 92-95.
3. Yang, B., et al. 1995. cDNA cloning, gene organization, and chromosomal localization of a human mercurial insensitive water channel. Evidence for distinct transcriptional units. *J. Biol. Chem.* 270: 22907-22913.
4. Ishibashi, K., et al. 1995. Structure and chromosomal localization of a human water channel (AQP3) gene. *Genomics* 27: 352-354.
5. Lee, M.D., et al. 1996. The human aquaporin-5 gene. Molecular characterization and chromosomal localization. *J. Biol. Chem.* 271: 8599-8604.
6. Turtzo, L.C., et al. 1997. Cloning and chromosomal localization of mouse aquaporin 4: exclusion of a candidate mutant phenotype, ataxia. *Genomics* 41: 267-270.
7. Frigeri, A., et al. 1998. Expression of aquaporin-4 in fast-twitch fibers of mammalian skeletal muscle. *J. Clin. Invest.* 102: 695-703.
8. Echevarria, M., et al. 1998. Aquaporins. *J. Physiol. Biochem.* 54: 107-118.
9. Beitz, E., et al. 1999. The mammalian aquaporin water channel family: A promising new drug target. *Curr. Med. Chem.* 6: 457-467.

CHROMOSOMAL LOCATION

Genetic locus: *Aqp4* (mouse) mapping to 18 A1.

PRODUCT

AQP4 (m): 293T Lysate represents a lysate of mouse AQP4 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

AQP4 (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive AQP4 antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.