



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

BHMT (m): 293T Lysate: sc-118804

BACKGROUND

Betaine-homocysteine methyltransferase (BHMT) is a zinc-dependent cytosolic protein that catalyzes the conversion of betaine and homocysteine to dimethylglycine and methionine, respectively. BHMT is highly expressed in rat liver, and its expression is regulated by dietary methionine and choline. In humans, S-adenosylmethionine (SAM) downregulates BHMT expression by inducing NFκB, which acts as a repressor for the BHMT gene. Lowered BHMT levels can lead to ER (endoplasmic reticulum) stress. Mutations in the gene encoding for BHMT may lead to hyperhomocysteinemia, a medical condition characterized by abnormally large amounts of homocysteine in the blood which may be a risk factor for cardiovascular and cerebrovascular diseases.

REFERENCES

1. Park, E.I. and Garrow, T.A. 1999. Interaction between dietary methionine and methyl donor intake on rat liver betaine-homocysteine methyltransferase gene expression and organization of the human gene. *J. Biol. Chem.* 274: 7816-7824.
2. Garrow, T.A. 2002. Random mutagenesis of the zinc-binding motif of betaine-homocysteine methyltransferase reveals that Gly 214 is essential. *Arch. Biochem. Biophys.* 399: 73-80.
3. Evans, J.C., Huddler, D.P., Jiracek, J., Castro, C., Millian, N.S., Garrow, T.A. and Ludwig, M.L. 2002. Betaine-homocysteine methyltransferase: zinc in a distorted barrel. *Structure* 10: 1159-1171.
4. Forestier, M., Bänninger, R., Reichen, J. and Solioz, M. 2003. Betaine homocysteine methyltransferase: gene cloning and expression analysis in rat liver cirrhosis. *Biochim. Biophys. Acta* 1638: 29-34.
5. Weisberg, I.S., Park, E., Ballman, K.V., Berger, P., Nunn, M., Suh, D.S., Garrow, T.A. and Rozen, R. 2003. Investigations of a common genetic variant in betaine-homocysteine methyltransferase (BHMT) in coronary artery disease. *Atherosclerosis* 167: 205-214.
6. Lee, M.B., Blunt, J.W., Lever, M. and George, P.M. 2004. A nuclear-magnetic-resonance-based assay for betaine-homocysteine methyltransferase activity. *Anal. Biochem.* 330: 199-205.
7. Castro, C., Gratson, A.A., Evans, J.C., Jiracek, J., Collinsová, M., Ludwig, M.L. and Garrow, T.A. 2004. Dissecting the catalytic mechanism of betaine-homocysteine S-methyltransferase by use of intrinsic tryptophan fluorescence and site-directed mutagenesis. *Biochemistry* 43: 5341-5351.
8. Zhu, H., Curry, S., Wen, S., Wicker, N.J., Shaw, G.M., Lammer, E.J., Yang, W., Jafarov, T. and Finnell, R.H. 2005. Are the betaine-homocysteine methyltransferase (BHMT and BHMT2) genes risk factors for spina bifida and orofacial clefts? *Am. J. Med. Genet. A* 135: 274-277.
9. Ou, X., Yang, H., Ramani, K., Ara, A.I., Chen, H., Mato, J.M. and Lu, S.C. 2006. Inhibition of human betaine-homocysteine methyltransferase expression by S-adenosylmethionine and methylthioadenosine. *Biochem. J.* 401: 87-96.

CHROMOSOMAL LOCATION

Genetic locus: Bhmt (mouse) mapping to 13 C3.

PRODUCT

BHMT (m): 293T Lysate represents a lysate of mouse BHMT transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.


APPLICATIONS

BHMT (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive BHMT antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

BHMT (3D6): sc-69708 is recommended as a positive control antibody for Western Blot analysis of enhanced mouse BHMT expression in BHMT transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

DATA



BHMT (3D6): sc-69708. Western blot analysis of BHMT expression in non-transfected: sc-117752 (A) and mouse BHMT transfected: sc-118804 (B) 293T whole cell lysates.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.