



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



CacyBP (m): 293T Lysate: sc-118948

BACKGROUND

CacyBP (calcyclin-binding protein, SIAH-interacting protein) is a 228 amino acid protein encoded by the human gene CACYBP. CacyBP is primarily a nuclear protein that contains one CS domain and one SGS domain. CacyBP is believed to be involved in calcium-dependent ubiquitination and subsequent proteosomal degradation of target proteins. It most likely serves as a molecular bridge in ubiquitin E3 complexes. It also participates in the ubiquitin-mediated degradation of β -catenin. CacyBP is thought to be a potential inhibitor of cell growth and invasion in the gastric cancer cell through its effects on β -catenin protein expression and transcriptional activation of TCF/LEF.

REFERENCES

1. Hildebrandt, B., Schoeler, D., Ringel, F., Kerner, T., Wust, P., Riess, H. and Schriever, F. 2007. Differential gene expression in peripheral blood lymphocytes of cancer patients treated with whole body hyperthermia and chemotherapy: a pilot study. 22: 625-635.
2. Sun, S., Ning, X., Liu, J., Liu, L., Chen, Y., Han, S., Zhang, Y., Liang, J., Wu, K. and Fan, D. 2007. Overexpressed CacyBP/SIP leads to the suppression of growth in renal cell carcinoma. Biochem. Biophys. Res. Commun. 356: 864-871.
3. Mei, Y., Xie, C., Xie, W., Wu, Z. and Wu, M. 2007. SIAH-1S, a novel splice variant of SIAH-1 (seven in absentia homolog), counteracts SIAH-1-mediated downregulation of β -catenin. Oncogene 26: 6319-6331.
4. Herington, J.L., Bi, J., Martin, J.D. and Bany, B.M. 2007. β -catenin (CTNNB) in the mouse uterus during decidualization and the potential role of two pathways in regulating its degradation. J. Histochem. Cytochem. 55: 963-974.
5. Liang, J., Luo, G., Ning, X., Shi, Y., Zhai, H., Sun, S., Jin, H., Liu, Z., Zhang, F., Lu, Y., Zhao, Y., Chen, X., Zhang, H., Guo, X., Wu, K. and Fan, D. 2007. Differential expression of calcium-related genes in gastric cancer cells transfected with cellular prion protein. Biochem. Cell Biol. 85: 375-383.
6. Schneider, G., Nieznanski, K., Kilanczyk, E., Bieganowski, P., Kuznicki, J. and Filipek, A. 2007. CacyBP/SIP interacts with Tubulin in neuroblastoma NB2a cells and induces formation of globular Tubulin assemblies. Biochim. Biophys. Acta 1773: 1628-1636.
7. Ning, X., Sun, S., Hong, L., Liang, J., Liu, L., Han, S., Liu, Z., Shi, Y., Li, Y., Gong, W., Zhang, S., Chen, Y., Guo, X., Cheng, Y., Wu, K. and Fan, D. 2008. Calcyclin-binding protein inhibits proliferation, tumorigenicity, and invasion of gastric cancer. Mol. Cancer Res. 5: 1254-1262.

CHROMOSOMAL LOCATION

Genetic locus: Cacybp (mouse) mapping to 1 H2.1.

PRODUCT

CacyBP (m): 293T Lysate represents a lysate of mouse CacyBP transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

CacyBP (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive CacyBP antibodies. Recommended use: 10-20 μ l per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.