



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



CARP (m): 293T Lysate: sc-119002

BACKGROUND

A proposed genetic marker of cardiac hypertrophy, CARP (cardiac ankyrin repeat protein) is a nuclear protein with an established role in regulation of cardiac gene expression. A distinct increase in CARP expression occurs in rats with abdominal aorta constriction, spontaneous hypertension and Dahl salt-sensitivity. In cardiomyocytes, CARP inhibits transcription of both cardiac Troponin C and atrial natriuretic factor. Specifically, expression of the CARP gene, which lies downstream of the cardiac homeobox gene Nkx-2.5, inhibits Nkx-2.5 transactivation of atrial natriuretic factor promoter. An increase in CARP expression is observed in the ventricular tissue of patients with end-stage heart failure. The major Ca^{2+} binding protein of cardiac sarcoplasmic reticulum (SR), calsequestrin (CSQ), upregulates the CARP gene and may contribute to the development of cardiac hypertrophy and fibrosis. $\text{TGF}\beta$ induces CARP expression in vascular smooth muscle cells (VSMCs), wherein CARP may mediate the inhibitory effects of $\text{TGF}\beta$ on VSMC proliferation.

REFERENCES

1. Jeyaseelan, R., Poizat, C., Baker, R.K., Abdishoo, S., Isterabadi, L.B., Lyons, G.E. and Kedes, L. 1997. A novel cardiac-restricted target for doxorubicin. CARP, a nuclear modulator of gene expression in cardiac progenitor cells and cardiomyocytes. *J. Biol. Chem.* 272: 22800-22808.
2. Zou, Y., Evans, S., Chen, J., Kuo, H.C., Harvey, R.P. and Chien, K.R. 1997. CARP, a cardiac ankyrin repeat protein, is downstream in the Nkx-2.5 homeobox gene pathway. *Development* 124: 793-804.
3. Aihara, Y., Kurabayashi, M., Saito, Y., Ohyama, Y., Tanaka, T., Takeda, S., Tomaru, K., Sekiguchi, K., Arai, M., Nakamura, T. and Nagai, R. 2000. Cardiac ankyrin repeat protein is a novel marker of cardiac hypertrophy: role of MCAT element within the promoter. *Hypertension* 36: 48-53.
4. Kanai, H., Tanaka, T., Aihara, Y., Takeda, S., Kawabata, M., Miyazono, K., Nagai, R. and Kurabayashi, M. 2001. Transforming growth factor- β /Smads signaling induces transcription of the cell type-restricted ankyrin repeat protein CARP gene through CagA motif in vascular smooth muscle cells. *Circ. Res.* 88: 30-36.
5. Bang, M.L., Mudry, R.E., McElhinny, A.S., Trombitas, K., Geach, A.J., Yamasaki, R., Sorimachi, H., Granzier, H., Gregorio, C.C. and Labeit, S. 2001. Myopalladin, a novel 145-kilodalton sarcomeric protein with multiple roles in Z-disc and I-band protein assemblies. *J. Cell Biol.* 153: 413-427.
6. Zolk, O., Frohme, M., Maurer, A., Kluxen, F.W., Hentsch, B., Zubakov, D., Hoheisel, J.D., Zucker, I.H., Pepe, S. and Eschenhagen, T. 2002. Cardiac ankyrin repeat protein, a negative regulator of cardiac gene expression, is augmented in human heart failure. *Biochem. Biophys. Res. Commun.* 293: 1377-1382.
7. Ihara, Y., Suzuki, Y.J., Kitta, K., Jones, L.R. and Ikeda, T. 2002. Modulation of gene expression in transgenic mouse hearts overexpressing calsequestrin. *Cell Calcium* 32: 21-29.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

CHROMOSOMAL LOCATION

Genetic locus: Ankrd1 (mouse) mapping to 19 C2.

PRODUCT

CARP (m): 293T Lysate represents a lysate of mouse CARP transfected 293T cells and is provided as 100 μg protein in 200 μl SDS-PAGE buffer.

APPLICATIONS

CARP (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive CARP antibodies. Recommended use: 10-20 μl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.