



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



CD164 (m): 293T Lysate: sc-119096

BACKGROUND

CD164 is a mucin-like cell surface glycoprotein that facilitates adhesion of CD34⁺ cells and serves as a negative regulator of hematopoietic progenitor cell proliferation. Human CD164 in CD34⁺/CD38⁺ hematopoietic progenitor and epithelial cell lines localizes to endosomes and lysosomes, with low concentrations also appearing at the cell surface.

REFERENCES

1. Watt, S.M., Bühring, H.J., Rappold, I., Chan, J.Y., Lee-Prudhoe, J., Jones, T., Zannettino, A.C., Simmons, P.J., Doyonnas, R., Sheer, D. and Butler, L.H. 1998. CD164, a novel sialomucin on CD34⁺ and erythroid subsets, is located on human chromosome 6q21. *Blood* 92: 849-866.
2. Doyonnas, R., Yi-Hsin Chan, J., Butler, L.H., Rappold, I., Lee-Prudhoe, J.E., Zannettino, A.C., Simmons, P.J., Bühring, H.J., Levesque, J.P. and Watt, S.M. 2000. CD164 monoclonal antibodies that block hematopoietic progenitor cell adhesion and proliferation interact with the first mucin domain of the CD164 receptor. *J. Immunol.* 165: 840-851.
3. Watt, S.M., Butler, L.H., Tavian, M., Bühring, H.J., Rappold, I., Simmons, P.J., Zannettino, A.C., Buck, D., Fuchs, A., Doyonnas, R., Chan, J.Y., Levesque, J.P., Peault, B. and Roxanis, I. 2000. Functionally defined CD164 epitopes are expressed on CD34⁺ cells throughout ontogeny but display distinct distribution patterns in adult hematopoietic and nonhematopoietic tissues. *Blood* 95: 3113-3124.
4. Chan, J.Y., Lee-Prudhoe, J.E., Jorgensen, B., Ihrke, G., Doyonnas, R., Zannettino, A.C., Buckle, V.J., Ward, C.J., Simmons, P.J. and Watt, S.M. 2001. Relationship between novel isoforms, functionally important domains, and subcellular distribution of CD164/endolyn. *J. Biol. Chem.* 276: 2139-2152.
5. Lee, Y.N., Kang, J.S. and Krauss, R.S. 2001. Identification of a role for the sialomucin CD164 in myogenic differentiation by signal sequence trapping in yeast. *Mol. Cell. Biol.* 21: 7696-7706.
6. McGuckin, C.P., Forraz, N., Baradez, M.O., Lojo-Rial, C., Wertheim, D., Whiting, K., Watt, S.M. and Pettengell, R. 2003. Colocalization analysis of sialomucins CD34 and CD164. *Stem Cells* 21: 162-170.
7. Jorgensen-Tye, B., Levesque, J.P., Royle, L., Doyonnas, R., Chan, J.Y., Dwek, R.A., Rudd, P.M., Harvey, D.J., Simmons, P.J. and Watt, S.M. 2005. Epitope recognition of antibodies that define the sialomucin, endolyn (CD164), a negative regulator of haematopoiesis. *Tissue Antigens* 65: 220-239.
8. Havens, A.M., Jung, Y., Sun, Y.X., Wang, J., Shah, R.B., Bühring, H.J., Pienta, K.J. and Taichman, R.S. 2006. The role of sialomucin CD164 (MGC-24v or endolyn) in prostate cancer metastasis. *BMC Cancer* 6: 195.
9. Bae, G.U., Gaio, U., Yang, Y.J., Lee, H.J., Kang, J.S. and Krauss, R.S. 2008. Regulation of myoblast motility and fusion by the CXCR4-associated sialomucin, CD164. *J. Biol. Chem.* 283: 8301-8309.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

CHROMOSOMAL LOCATION

Genetic locus: Cd164 (mouse) mapping to 10 B2.

PRODUCT

CD164 (m): 293T Lysate represents a lysate of mouse CD164 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

APPLICATIONS

CD164 (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive CD164 antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.