



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

CD4 (m): 293T Lysate: sc-119107

BACKGROUND

The T cell receptor (TCR) is a heterodimer composed of either α and β or γ and δ chains. CD3 chains and the CD4 or CD8 co-receptors are also required for efficient signal transduction through the TCR. The TCR is expressed on T helper and T cytotoxic cells that can be distinguished by their expression of CD4 and CD8; T helper cells express CD4 proteins and T cytotoxic cells display CD8. CD4 is also expressed on cortical cells, mature medullary thymocytes, microglial cells and dendritic cells. CD4 (also designated T4 and Leu 3) is a membrane glycoprotein that contains four extracellular immunoglobulin-like domains. The TCR in association with CD4 can bind class II MHC molecules presented by the antigen-presenting cells. The CD4 protein functions by increasing the avidity of the interaction between the TCR and an antigen-class II MHC complex. An additional role of CD4 is to function as a receptor for HIV.

REFERENCES

- Maddon, P.J., et al. 1987. Structure and expression of human and mouse T4 genes. *Proc. Natl. Acad. Sci. USA* 84: 9155-9159.
- Arthos, J., et al. 1989. Identification of the residues in human CD4 critical for the binding of HIV. *Cell* 57: 469-481.
- Healey, D., et al. 1990. Novel anti-CD4 monoclonal antibodies separate human immunodeficiency virus infection and fusion of CD4⁺ cells from virus binding. *J. Exp. Med.* 172: 1233-1242.
- Allison, J.P., et al. 1991. The immunobiology of T cells with invariant γ δ antigen receptors. *Annu. Rev. Immunol.* 9: 679-705.
- Janeway, C.A., Jr. 1992. The T cell receptor as a multicomponent signalling machine: CD4/CD8 co-receptors and CD45 in T cell activation. *Annu. Rev. Immunol.* 10: 645-674.
- Ehrlich, E.W., et al. 1993. T cell receptor interaction with peptide/major histocompatibility complex (MHC) and superantigen/MHC ligands is dominated by antigen. *J. Exp. Med.* 178: 713-722.
- Julius, M., et al. 1993. Distinct roles for CD4 and CD8 as co-receptors in antigen receptor signalling. *Immunol. Today* 14: 177-183.
- Vignali, D.A. 1994. The interaction between CD4 and MHC class II molecules and its effect on T cell function. *Behring Inst. Mitt.* 94: 133-147.
- Schlossman, S., et al, eds. 1995. *Leucocyte Typing V*. Oxford: Oxford University Press.

CHROMOSOMAL LOCATION

Genetic locus: Cd4 (mouse) mapping to 6 F2.

PRODUCT

CD4 (m): 293T Lysate represents a lysate of mouse CD4 transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.


APPLICATIONS

CD4 (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive CD4 antibodies. Recommended use: 10-20 μ l per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

CD4 (RM4-5): sc-19643 is recommended as a positive control antibody for Western Blot analysis of enhanced mouse CD4 expression in CD4 transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

DATA



CD4 (RM4-5): sc-19643. Western blot analysis of CD4 expression in non-transfected: sc-117752 (A) and mouse CD4 transfected: sc-119107 (B) 293T whole cell lysates.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.