



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



Chx10 (m2): 293T Lysate: sc-119257

BACKGROUND

Chx10, for *ceh-10* homeodomain containing homolog, is also known as RET1 and HOX10 and is closely related to the homeodomain of the homeobox gene *ceh-10* from the nematode *Caenorhabditis elegans*. Chx10 is an essential component in the network of genes required for the development of the mammalian eye, with profound effects on retinal progenitor proliferation and bipolar cell specification or differentiation. Chx10 is expressed in the early retinal neuroepithelium, is restricted to bipolar cells and is maintained at a low level in bipolar cells of the mature retina. Human Chx10 is also expressed in the inner nuclear layer of the mature retina. Expression patterns implicate critical roles in the formation of the neuroretina and in the development and maintenance of the inner nuclear layer. Chx10 is expressed at high levels in uncommitted retinal progenitor cells and mature bipolar cells.

REFERENCES

1. Liu, I.S., Chen, J.D., Ploder, L., Vidgen, D., van der Kooy, D., Kalnins, V.I. and McInnes, R.R. 1994. Developmental expression of a novel murine homeobox gene (Chx10): evidence for roles in determination of the neuroretina and inner nuclear layer. *Neuron* 13: 377-393.
2. Svendsen, P.C. and McGhee, J.D. 1995. The *C. elegans* neuronally expressed homeobox gene *ceh-10* is closely related to genes expressed in the vertebrate eye. *Development* 121: 1253-1262.
3. Burmeister, M., Novak, J., Liang, M.Y., Basu, S., Ploder, L., Hawes, N.L., Vidgen, D., Hoover, F., Goldman, D., Kalnins, V.I., Roderick, T.H., Taylor, B.A., Hankin, M.H. and McInnes, R.R. 1996. Ocular retardation mouse caused by Chx10 homeobox null allele: impaired retinal progenitor proliferation and bipolar cell differentiation. *Nat. Genet.* 12: 376-384.
4. Chen, C.M. and Cepko, C.L. 2000. Expression of Chx10 and Chx10-1 in the developing chicken retina. *Mech. Dev.* 90: 293-297.
5. Ferda Percin, E., Ploder, L.A., Yu, J.J., Arici, K., Horsford, D.J., Rutherford, A., Bapat, B., Cox, D.W., Duncan, A.M., Kalnins, V.I., Kocak-Altintas, A., Sowden, J.C., Traboulsi, E., Sarfarazi, M. and McInnes, R.R. 2000. Human microphthalmia associated with mutations in the retinal homeobox gene Chx10. *Nat. Genet.* 25: 397-401.
6. Online Mendelian Inheritance in Man, OMIM™. 2000. Johns Hopkins University, Baltimore, MD. MIM Number: 142993. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
7. LocusLink Report (LocusID: 1148). <http://www.ncbi.nlm.nih.gov/LocusLink/>

CHROMOSOMAL LOCATION

Genetic locus: *Vsx2* (mouse) mapping to 12 D1.

PRODUCT

Chx10 (m2): 293T Lysate represents a lysate of mouse Chx10 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

APPLICATIONS

Chx10 (m2): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive Chx10 antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.