



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

CNOT10 (m2): 293T Lysate: sc-119340

BACKGROUND

CNOT10 is a subunit of the CCR4-NOT complex which consists of at least five other CNOT subunit proteins and TAB182. The CCR4-NOT complex is an evolutionarily conserved, multi-component complex known to be involved in transcription, as well as in mRNA degradation. Various subunits (e.g. CNOT1, CNOT3) are uniquely involved in influencing nuclear hormone receptor activities. In effect, this complex has an important role as a transcription regulator and repressor of nuclear receptor signaling that is relevant to the molecular pathways involved in cancer. The CCR4-NOT complex is also involved in the regulation of Histone H3 lysine 4 methylation through a ubiquitin-dependent pathway that likely involves the proteasome.

REFERENCES

- Mulder, K.W., Winkler, G.S. and Timmers, H.T. 2005. DNA damage and replication stress induced transcription of RNR genes is dependent on the CCR4-NOT complex. *Nucleic Acids Res.* 33: 6384-6392.
- Winkler, G.S., Mulder, K.W., Bardwell, V.J., Kalkhoven, E. and Timmers, H.T. 2006. Human CCR4-NOT complex is a ligand-dependent repressor of nuclear receptor-mediated transcription. *EMBO J.* 25: 3089-3099.
- Behm-Ansmant, I., Rehwinkel, J., Doerks, T., Stark, A., Bork, P. and Izaurralde, E. 2006. mRNA degradation by miRNAs and GW182 requires both CCR4:NOT deadenylase and DCP1:DCP2 decapping complexes. *Genes Dev.* 20: 1885-1898.
- Lenssen, E., Azzouz, N., Michel, A., Landrieux, E. and Collart, M.A. 2007. The CCR4-NOT complex regulates Skn7 through Srb10 kinase. *Eukaryotic Cell* 6: 2251-2259.
- James, N., Landrieux, E. and Collart, M.A. 2007. A SAGA-independent function of SPT3 mediates transcriptional deregulation in a mutant of the CCR4-NOT complex in *Saccharomyces cerevisiae*. *Genetics* 177: 123-135.
- Wagner, E., Clement, S.L. and Lykke-Andersen, J. 2007. An unconventional human CCR4-CAF1 deadenylase complex in nuclear cajal bodies. *Mol. Cell Biol.* 27: 1686-1695.
- Mulder, K.W., Brenkman, A.B., Inagaki, A., van den Broek, N.J. and Timmers, H.T. 2007. Regulation of histone H3K4 tri-methylation and PAF complex recruitment by the CCR4-NOT complex. *Nucleic Acids Res.* 35: 2428-2439.
- Larabee, R.N., Shibata, Y., Mersman, D.P., Collins, S.R., Kemmeren, P., Roguev, A., Weissman, J.S., Briggs, S.D., Krogan, N.J. and Strahl, B.D. 2007. CCR4/NOT complex associates with the proteasome and regulates histone methylation. *Proc. Natl. Acad. Sci. USA* 104: 5836-5841.
- Garapaty, S., Mahajan, M.A. and Samuels, H.H. 2008. Components of the CCR4-NOT complex function as nuclear hormone receptor coactivators via association with the NRC-interacting factor NIF-1. *J. Biol. Chem.* 283: 6806-6816.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

CHROMOSOMAL LOCATION

Genetic locus: Cnot10 (mouse) mapping to 9 F3.

PRODUCT

CNOT10 (m2): 293T Lysate represents a lysate of mouse CNOT10 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

APPLICATIONS

CNOT10 (m2): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive CNOT10 antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.