



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

CRISP-3 (m): 293T Lysate: sc-119466

BACKGROUND

Cysteine-rich secretory proteins (CRISPs) represent a family of evolutionarily conserved proteins which may play a role in the innate immune system and are transcriptionally regulated by androgens in several tissues. AEG is a sperm surface protein involved in the fusion of egg and sperm. Although CRISP-1 (also designated AEG-like protein, ARP, cysteine-rich secretory protein-1 or AEG-related protein) is not the ortholog of rodent AEG, it resembles AEG in that it is an epididymal secretory glycoprotein that binds to the postacrosomal region of the sperm head. CRISP-1 coats the postacrosomal region of sperm heads as they pass through the epididymis. CRISP-1 is found in all regions of the epididymis, ductus deferens, seminal plasma and sperm. CRISP-3 is expressed in pancreas and prostate tissues and, along with CRISP-1, is expressed in saliva. The gene that encodes CRISP-3 is an early response gene that may participate in the pathophysiology of the autoimmune lesions of Sjogren's syndrome.

REFERENCES

1. Krätzschar, J., et al. 1996. The human cysteine-rich secretory protein (CRISP) family. Primary structure and tissue distribution of CRISP-1, CRISP-2 and CRISP-3. *Eur. J. Biochem.* 236: 827-836.
2. Liao, Q., et al. 2003. Preferential expression of cystein-rich secretory protein-3 (CRISP-3) in chronic pancreatitis. *Histol. Histopathol.* 18: 425-433.
3. Udby, L., et al. 2004. Cysteine-rich secretory protein 3 is a ligand of α 1B-glycoprotein in human plasma. *Biochemistry* 43: 12877-12886.
4. Udby, L., et al. 2005. β -Microseminoprotein binds CRISP-3 in human seminal plasma. *Biochem. Biophys. Res. Commun.* 333: 555-561.
5. Udby, L., et al. 2005. Characterization and localization of cysteine-rich secretory protein 3 (CRISP-3) in the human male reproductive tract. *J. Androl.* 26: 333-342.
6. Bjartell, A., et al. 2006. Immunohistochemical detection of cysteine-rich secretory protein 3 in tissue and in serum from men with cancer or benign enlargement of the prostate gland. *Prostate* 66: 591-603.
7. Laine, M., et al. 2007. Low salivary dehydroepiandrosterone and androgen-regulated cysteine-rich secretory protein 3 levels in Sjögren's syndrome. *Arthritis Rheum.* 56: 2575-2584.
8. Bjartell, A.S., et al. 2007. Association of cysteine-rich secretory protein 3 and β -Microseminoprotein with outcome after radical prostatectomy. *Clin. Cancer Res.* 13: 4130-4138.
9. Ghasriani, H., et al. 2009. A model of the complex between human β -Microseminoprotein and CRISP-3 based on NMR data. *Biochem. Biophys. Res. Commun.* 378: 235-239.

CHROMOSOMAL LOCATION

Genetic locus: Crisp3 (mouse) mapping to 17 B2.

PRODUCT

CRISP-3 (m): 293T Lysate represents a lysate of mouse CRISP-3 transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

APPLICATIONS

CRISP-3 (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive CRISP-3 antibodies. Recommended use: 10-20 μ l per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.