



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

CYB561D2 (m): 293T Lysate: sc-119539

BACKGROUND

CYB561D2 (cytochrome b561 domain-containing protein 2), also known as putative tumor suppressor protein 101F6, is a 222 amino acid highly hydrophobic protein that contains 6 membrane spanning domains. Localized to endosomes and endoplasmic reticulum of the perinuclear region, CYB561D2 contains 2 heme-binding sites, through which it reduces ferric ions. CYB561D2 is ubiquitously expressed, with highest levels in lung, liver and kidney. Though expression of CYB561D2 is abundant in normal lung bronchial epithelial cells, CYB561D2 is absent in most lung cancers. The gene encoding CYB561D2 maps to human chromosome 3p21.31, a region that is prone to genetic alterations and is associated with frequent and early allele loss in many human cancers. Exogenous expression of CYB561D2 leads to the induction of caspase-independent apoptotic and autophagic pathways, resulting in the synergistic killing of tumor cells. This suggests an important role for CYB561D2 as a therapeutic intervention in cancer treatment.

REFERENCES

- Lerman, M.I. and Minna, J.D. 2000. The 630-kb lung cancer homozygous deletion region on human chromosome 3p21.3: identification and evaluation of the resident candidate tumor suppressor genes. The International Lung Cancer Chromosome 3p21.3 Tumor Suppressor Gene Consortium. *Cancer Res.* 60: 6116-6133.
- Ji, L., Nishizaki, M., Gao, B., Burbee, D., Kondo, M., Kamibayashi, C., Xu, K., Yen, N., Atkinson, E.N., Fang, B., Lerman, M.I., Roth, J.A. and Minna, J.D. 2002. Expression of several genes in the human chromosome 3p21.3 homozygous deletion region by an adenovirus vector results in tumor suppressor activities *in vitro* and *in vivo*. *Cancer Res.* 62: 2715-2720.
- Chow, L.S., Lo, K.W., Kwong, J., To, K.F., Tsang, K.S., Lam, C.W., Dammann, R. and Huang, D.P. 2004. RASSF1A is a target tumor suppressor from 3p21.3 in nasopharyngeal carcinoma. *Int. J. Cancer* 109: 839-847.
- Angeloni, D. 2007. Molecular analysis of deletions in human chromosome 3p21 and the role of resident cancer genes in disease. *Brief. Funct. Genomic Proteomic* 6: 19-39.
- Ohtani, S., Iwamaru, A., Deng, W., Ueda, K., Wu, G., Jayachandran, G., Kondo, S., Atkinson, E.N., Minna, J.D., Roth, J.A. and Ji, L. 2007. Tumor suppressor 101F6 and ascorbate synergistically and selectively inhibit non-small cell lung cancer growth by caspase-independent apoptosis and autophagy. *Cancer Res.* 67: 6293-6303.
- Mizutani, A., Sanuki, R., Kakimoto, K., Kojo, S. and Taketani, S. 2007. Involvement of 101F6, a homologue of cytochrome b561, in the reduction of ferric ions. *J. Biochem.* 142: 699-705.
- Hesson, L.B., Cooper, W.N. and Latif, F. 2007. Evaluation of the 3p21.3 tumour-suppressor gene cluster. *Oncogene* 26: 7283-7301.
- Online Mendelian Inheritance in Man, OMIM[™]. 2008. Johns Hopkins University, Baltimore, MD. MIM Number: 607068. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>

CHROMOSOMAL LOCATION

Genetic locus: Cyb561d2 (mouse) mapping to 9 F1.

PRODUCT

CYB561D2 (m): 293T Lysate represents a lysate of mouse CYB561D2 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

APPLICATIONS

CYB561D2 (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive CYB561D2 antibodies. Recommended use: 10-20 µl per lane.


Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

CYB561D2 (C-2): sc-373870 is recommended as a positive control antibody for Western Blot analysis of enhanced mouse CYB561D2 expression in CYB561D2 transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:
1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker[™] Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

DATA



CYB561D2 (C-2): sc-373870. Western blot analysis of CYB561D2 expression in non-transfected: sc-117752 (A) and mouse CYB561D2 transfected: sc-119539 (B) 293T whole cell lysates.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.