



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

DAP-1 (m2): 293T Lysate: sc-119658

BACKGROUND

In contrast to growth factors which promote cell proliferation, FAS ligand (FAS-L) and the tumor necrosis factors (TNFs) rapidly induce apoptosis. Cellular response to FAS-L and TNF is mediated by structurally related receptors containing a conserved cytoplasmic region called the "death domain". DAP-1 (for death associated protein-1) is a basic, proline-rich protein expressed in γ interferon (IFN- γ)-induced HeLa cells. DAP-1 is a member of the ubiquitin homology (UbH) family which also includes SUMO-1 and SIII p18 elogin protein. DAP-1 interacts with the death domain of TNF-R1 and can trigger programmed cell death in a variety of cell lines, as well as suppress NF κ B/Rel activity.

REFERENCES

1. Tartaglia, L.A., et al. 1993. A novel domain within the 55 kDa TNF receptor signals cell death. *Cell* 74: 845-853.
2. Itoh, N., et al. 1993. A novel protein domain required for apoptosis. Mutational analysis of human FAS antigen. *J. Biol. Chem.* 268: 10932-10937.
3. Smith, C.A., et al. 1994. The TNF receptor superfamily of cellular and viral proteins: activation, costimulation, and death. *Cell* 76: 959-962.
4. Nagata, S., et al. 1995. The FAS death factor. *Science* 267: 1449-1456.
5. Deiss, L.P., et al. 1995. Identification of a novel serine/threonine kinase and a novel 15 kDa protein as potential mediators of the γ interferon-induced cell death. *Genes Dev.* 9: 15-30.
6. Garrett, K.P., et al. 1995. Positive regulation of general transcription factor SIII by a tailed ubiquitin homolog. *Proc. Natl. Acad. Sci. USA* 92: 7172-7176.
7. Boddy, M.N., et al. 1996. PIC 1, a novel ubiquitin-like protein which interacts with the PML component of a multiprotein complex that is disrupted in acute promyelocytic leukemia. *Oncogene* 13: 971-982.
8. Ware, C.F., et al. 1996. Apoptosis mediated by the TNF-related cytokine and receptor families. *J. Cell. Biochem.* 60: 47-55.
9. Liou, M.L., et al. 1999. The ubiquitin-homology protein, DAP-1, associates with tumor necrosis factor receptor (p60) death domain and induces apoptosis. *Biol. Chem.* 274: 10145-10153.

CHROMOSOMAL LOCATION

Genetic locus: Dap (mouse) mapping to 15 B2.

PRODUCT

DAP-1 (m2): 293T Lysate represents a lysate of mouse DAP-1 transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

STORAGE

Store at -20 $^{\circ}$ C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

APPLICATIONS

DAP-1 (m2): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive DAP-1 antibodies. Recommended use: 10-20 μ l per lane.


Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

DAP-1 (C-8): sc-376754 is recommended as a positive control antibody for Western Blot analysis of enhanced mouse DAP-1 expression in DAP-1 transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:
 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz MarkerTM Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

DATA



DAP-1 (C-8): sc-376754. Western blot analysis of DAP-1 expression in non-transfected: sc-117752 (A) and mouse DAP-1 transfected: sc-119658 (B) 293T whole cell lysates.

RESEARCH USE

For research use only, not for use in diagnostic procedures.