



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

DHODH (m2): 293T Lysate: sc-119762

BACKGROUND

DHODH (dihydroorotate dehydrogenase), also known as DHodehase, is a 395 amino acid mitochondrial protein located on the outer surface of the inner mitochondrial membrane. It catalyzes the fourth enzymatic step in *de novo* pyrimidine biosynthesis. *De novo* pyrimidine synthesis is a critical metabolic pathway for nucleic acid synthesis and is a target for various cancer chemotherapy agents. Additionally, DHODH is functionally connected to the respiratory chain, delivering electrons to ubiquinone. DHODH contains a bipartite signal at the N-terminus that regulates passage into the mitochondrial inner membrane. The inhibition of COX (cytochrome c oxidase) by nitric oxide (NO) indirectly inhibits DHODH activity. The inhibition of DHODH has an immunosuppressive and an antiproliferative effect on diseases such as rheumatoid arthritis.

REFERENCES

1. Barnes, T., et al. 1993. Regional mapping of the gene encoding dihydroorotate dehydrogenase, an enzyme involved in UMP synthesis, electron transport, and superoxide generation, to human chromosome region 16q22. *Somat. Cell Mol. Genet.* 19: 405-411.
2. Copeland, R.A., et al. 1995. Recombinant human dihydroorotate dehydrogenase: expression, purification, and characterization of a catalytically functional truncated enzyme. *Arch. Biochem. Biophys.* 323: 79-86.
3. Knecht, W., et al. 1996. Functional expression of a fragment of human dihydroorotate dehydrogenase by means of the baculovirus expression vector system, and kinetic investigation of the purified recombinant enzyme. *Eur. J. Biochem.* 240: 292-301.
4. Beuneu, C., et al. 2000. Indirect inhibition of mitochondrial dihydroorotate dehydrogenase activity by nitric oxide. *Free Radic. Biol. Med.* 28: 1206-1213.
5. Dietz, C., et al. 2000. Immunocytochemical detection of mitochondrial dihydroorotate dehydrogenase in human spermatozoa. *Int. J. Androl.* 23: 294-299.
6. Rawls, J., et al. 2000. Requirements for the mitochondrial import and localization of dihydroorotate dehydrogenase. *Eur. J. Biochem.* 267: 2079-2087.
7. Small, Y.A., et al. 2006. Hydrogen bonding pathways in human dihydroorotate dehydrogenase. *J. Phys. Chem. B* 110: 19704-19710.
8. Baumgartner, R., et al. 2006. Dual binding mode of a novel series of DHODH inhibitors. *J. Med. Chem.* 49: 1239-1247.
9. Zameitat, E., et al. 2007. Functional expression of human dihydroorotate dehydrogenase (DHODH) in *pyr4* mutants of *Ustilago maydis* allows target validation of DHODH inhibitors *in vivo*. *Appl. Environ. Microbiol.* 73: 3371-3379.

CHROMOSOMAL LOCATION

Genetic locus: Dhodh (mouse) mapping to 8 D3.

PRODUCT

DHODH (m2): 293T Lysate represents a lysate of mouse DHODH transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

APPLICATIONS

DHODH (m2): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive DHODH antibodies. Recommended use: 10-20 µl per lane.


Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

DHODH (E-8): sc-166348 is recommended as a positive control antibody for Western Blot analysis of enhanced mouse DHODH expression in DHODH transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

DATA



DHODH (E-8): sc-166348. Western blot analysis of DHODH expression in non-transfected: sc-117752 (A) and mouse DHODH transfected: sc-119762 (B) 293T whole cell lysates.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.