



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



EHD (m): 293T Lysate: sc-119958

BACKGROUND

The Eps15 homology (EH) domain-containing protein family consists of four members, EHD1, EHD2, EHD3, and EHD4. The chromosomal locations of the human EHD genes are as follows: EHD1 maps to 11q13.1, EHD2 maps to 19q13.33, EHD3 maps to 2p23.1, and EHD4 maps to 15q15.1. The encoded proteins of all EHD family members contain multiple conserved regions, which include an amino-terminal nucleotide-binding consensus site, a bipartite nuclear localization signal, and a carboxy-terminal EH protein-binding domain with an EF-hand motif. EHD1 is ubiquitously expressed with increased expression in testis. EHD2, EHD3, and EHD4 have more specific expression with EHD2 highly expressed in heart, EHD3 expressed in brain, kidney, liver, placenta, ovary, and heart, and EHD4 expressed in heart, placenta, and pancreas. The EHD proteins may participate in ligand-induced endocytosis.

REFERENCES

- Haider, N.B., et al. 1999. Evaluation and molecular characterization of EHD1, a candidate gene for Bardet-Biedl syndrome 1 (BBS1). *Gene* 240: 227-232.
- Mintz, L., et al. 1999. EHD1—an EH-domain-containing protein with a specific expression pattern. *Genomics* 59: 66-76.
- Pohl, U., et al. 2000. EHD2, EHD3, and EHD4 encode novel members of a highly conserved family of EH domain-containing proteins. *Genomics* 63: 255-262.
- Kuo, H.J., et al. 2001. Characterization of EHD4, an EH domain-containing protein expressed in the extracellular matrix. *J. Biol. Chem.* 276: 43103-43110.
- Caplan, S., et al. 2002. A tubular EHD1-containing compartment involved in the recycling of major histocompatibility complex class I molecules to the plasma membrane. *EMBO J.* 21: 2557-2567.
- Galperin, E., et al. 2002. EHD3: a protein that resides in recycling tubular and vesicular membrane structures and interacts with EHD1. *Traffic* 3: 575-589.

PRODUCT

EHD (m): 293T Lysate represents a lysate of mouse EHD transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

EHD (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive EHD antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

EHD (E-5): sc-271658 is recommended as a positive control antibody for Western Blot analysis of enhanced mouse EHD expression in EHD transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:

- Western Blotting: use m-IgG_X BP-HRP: sc-516102 or m-IgG_X BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

DATA



EHD (E-5): sc-271658. Western blot analysis of EHD expression in non-transfected: sc-117752 (**A**) and mouse EHD transfected: sc-119958 (**B**) 293T whole cell lysates.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.