



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# ENX-1 (m): 293T Lysate: sc-120052

## BACKGROUND

In *Drosophila*, the Polycomb (PcG) gene family encodes chromatin proteins that are required for the repression of homeotic loci in embryonic development. PcG proteins work in conjunction with the trithorax group (trxG), which activate homeobox gene expression during embryonic development. ENX-1, a mammalian homolog of the *Drosophila* gene enhancer of zeste, is a PcG protein that is ubiquitously expressed during early embryogenesis and becomes restricted to the central and peripheral nervous systems and sites of fetal hematopoiesis during later development. In the adult, ENX-1 is restricted to specific sites, including spleen, testis and placenta. The gene encoding human ENX-1 transcribes a 746 amino acid polypeptide which contains a trithorax-like domain and a DNA-binding motif. ENX-1 interacts with the proto-oncogene product Vav and is thought to be involved in the proliferation of normal and malignant hematopoietic cells. By altering the regulation of target genes, ENX-1 may also contribute to certain phenotypes of Down syndrome.

## REFERENCES

1. Goebel, M.G. 1991. The Bmi-1 and Mel-18 gene products define a new family of DNA-binding proteins involved in cell proliferation and tumorigenesis. *Cell* 66: 623.
2. Hobert, O., et al. 1996. Isolation and developmental expression analysis of ENX-1, a novel mouse Polycomb group gene. *Mech. Dev.* 55: 171-184.
3. Hobert, O., et al. 1996. Interaction of Vav with ENX-1, a putative transcriptional regulator of homeobox gene expression. *Mol. Cell. Biol.* 16: 3066-3073.
4. Laible, G., et al. 1997. Mammalian homologues of the Polycomb group gene enhancer of zeste mediate gene silencing in *Drosophila* heterochromatin and at *S. cerevisiae* telomeres. *EMBO J.* 16: 3219-3232.
5. van Lohuizen, M., et al. 1998. Interaction of mouse polycomb group (PcG) proteins ENX-1 and ENX-2 with EED: indication for separate PcG complexes. *Mol. Cell. Biol.* 18: 3572-3579.
6. Sewalt, R.G., et al. 1998. Characterization of interactions between the mammalian polycomb group proteins ENX-1/EZH2 and EED suggests the existence of different mammalian Polycomb-group protein complexes. *Mol. Cell. Biol.* 18: 3586-3595.
7. Fukuyama, T., et al. 2000. Proliferative involvement of ENX-1, a putative human polycomb group gene, in haematopoietic cells. *Br. J. Haematol.* 108: 842-847.

## CHROMOSOMAL LOCATION

Genetic locus: Ezh2 (mouse) mapping to 6 B2.3.

## PRODUCT

ENX-1 (m): 293T Lysate represents a lysate of mouse ENX-1 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## APPLICATIONS

ENX-1 (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive ENX-1 antibodies. Recommended use: 10-20 µl per lane.


Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

ENX-1 (C-1): sc-166609 is recommended as a positive control antibody for Western Blot analysis of enhanced mouse ENX-1 expression in ENX-1 transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

## RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:  
 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

## DATA



ENX-1 (C-1): sc-166609. Western blot analysis of ENX-1 expression in non-transfected: sc-117752 (A) and mouse ENX-1 transfected: sc-120052 (B) 293T whole cell lysates.

## STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.