



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

EPAS-1 (m): 293T Lysate: sc-120061

BACKGROUND

Cell growth and viability is compromised by oxygen deprivation (hypoxia). Hypoxia-inducible factors, including HIF-1 α , HIF-1 β (also designated Arnt 1), EPAS-1 (also designated HIF-2 α) and HIF-3 α , induce glycolysis, erythropoiesis and angiogenesis in order to restore oxygen homeostasis. Hypoxia-inducible factors are members of the Per-Arnt-Sim (PAS) domain transcription factor family. In response to hypoxia, HIF-1 α is upregulated and forms a heterodimer with Arnt 1 to form the HIF-1 complex. The HIF-1 complex recognizes and binds to the hypoxia responsive element (HRE) of hypoxia-inducible genes, thereby activating transcription. Hypoxia-inducible expression of some genes such as Glut-1, p53, p21 or Bcl-2, is HIF-1 α dependent, whereas expression of others, such as p27, GADD 153 or HO-1, is HIF-1 α independent. EPAS-1 and HIF-3 α have also been shown to form heterodimeric complexes with Arnt 1 in response to hypoxia.

REFERENCES

1. Wang, G.L., Jiang, B.H., Rue, E.A. and Semenza, G.L. 1995. Hypoxia-inducible factor 1 is a basic-helix-loop-helix-PAS heterodimer regulated by cellular O₂ tension. *Proc. Natl. Acad. Sci. USA* 92: 5510-5514.
2. Tian, H., McKnight, S.L. and Russell, D.W. 1997. Endothelial PAS domain protein-1 (EPAS-1), a transcription factor selectively expressed in endothelial cells. *Genes Dev.* 11: 72-82.
3. Luo, G., Gu, Y.Z., Jain, S., Chan, W.K., Carr, K.M., Hogenesch, J.B. and Bradfield, C.A. 1997. Molecular characterization of the murine Hif-1 α locus. *Gene Expr.* 6: 287-299.
4. Carmeliet, P., Dor, Y., Herbert, J.M., Fukumura, D., Brusselmans, K., Dewerchin, M., Neeman, M., Bono, F., Abramovitch, R., Maxwell, P., Koch, C.J., Ratcliffe, P., Moons, L., Jain, R.K., Collen, D. and Keshet, E. 1998. Role of HIF-1 α in hypoxia-mediated apoptosis, cell proliferation and tumour angiogenesis. *Nature* 394: 485-490.
5. Gu, Y.Z., Moran, S.M., Hogenesch, J.B., Wartman, L. and Bradfield, C.A. 1998. Molecular characterization and chromosomal localization of a third α -class hypoxia inducible factor subunit, HIF-3 α . *Gene Expr.* 7: 205-213.
6. Wood, S.M., Wiesener, M.S., Yeates, K.M., Okada, N., Pugh, C.W., Maxwell, P.H. and Ratcliffe, P.J. 1998. Selection and analysis of a mutant cell line defective in the hypoxia-inducible factor-1 α subunit (HIF-1 α). Characterization of HIF-1 α -dependent and -independent hypoxia-inducible gene expression. *J. Biol. Chem.* 273: 8360-8368.

CHROMOSOMAL LOCATION

Genetic locus: *Epas1* (mouse) mapping to 17 E4.

PRODUCT

EPAS-1 (m): 293T Lysate represents a lysate of mouse EPAS-1 transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

STORAGE

Store at -20 $^{\circ}$ C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

APPLICATIONS

EPAS-1 (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive EPAS-1 antibodies. Recommended use: 10-20 μ l per lane.


Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

EPAS-1 (190b): sc-13596 is recommended as a positive control antibody for Western Blot analysis of enhanced mouse EPAS-1 expression in EPAS-1 transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:
1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker[™] Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

DATA



EPAS-1 (190b): sc-13596. Western blot analysis of EPAS-1 expression in untreated (A) and CoCl₂-induced (B) HeLa whole cell lysate.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.