



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

GATAD1 (m): 293T Lysate: sc-120425

BACKGROUND

GATAD1 (GATA zinc finger domain-containing protein 1), also known as ODAG (ocular development-associated gene protein), is a 269 amino acid protein that is involved in early ocular development. Expressed highly in postnatal eye tissue, GATAD1 is associated with formation of the lens and its surrounding structures, suggesting a possible role in the transformation of ocular tissues into a working eye. GATAD1 expression declines dramatically after the early stages of development. GATAD1 contains 1 GATA-type zinc finger which functions as a DNA-binding domain. Additionally, GATAD1 gene expression is amplified in certain cancerous cells, suggesting that it may be involved in carcinogenesis.

REFERENCES

1. Tsuruga, T., Kanamoto, T., Kato, T., Yamashita, H., Miyagawa, K. and Mishima, H.K. 2002. Ocular development-associated gene (ODAG), a novel gene highly expressed in ocular development. *Gene* 290: 125-130.
2. Pang, E.Y., Bai, A.H., To, K.F., Sy, S.M., Wong, N.L., Lai, P.B., Squire, J.A. and Wong, N. 2007. Identification of PFTAIRE protein kinase 1, a novel cell division cycle-2 related gene, in the motile phenotype of hepatocellular carcinoma cells. *Hepatology* 46: 436-445.
3. Agelopoulos, K., Kersting, C., Korsching, E., Schmidt, H., Kuijper, A., August, C., Wülfing, P., Tio, J., Boecker, W., van Diest, P.J., Brandt, B. and Buerger, H. 2007. EGFR amplification specific gene expression in phyllodes tumours of the breast. *Cell. Oncol.* 29: 443-451.
4. Nagel, S., Leich, E., Quentmeier, H., Meyer, C., Kaufmann, M., Drexler, H.G., Zettl, A., Rosenwald, A. and Macleod, R.A. 2008. Amplification at 7q22 targets cyclin-dependent kinase 6 in T cell lymphoma. *Leukemia* 22: 387-392.

CHROMOSOMAL LOCATION

Genetic locus: Gatad1 (mouse) mapping to 5 A1.

PRODUCT

GATAD1 (m): 293T Lysate represents a lysate of mouse GATAD1 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

APPLICATIONS

GATAD1 (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive GATAD1 antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

RESEARCH USE

For research use only, not for use in diagnostic procedures.