



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# GP73 (m): 293T Lysate: sc-120573

## BACKGROUND

GP73 (also known as Golgi phosphoprotein 2, GOLPH 2 or Golgi membrane protein), is a widely expressed, epithelial-specific, type II transmembrane protein which resides in the Golgi apparatus, where it is responsible for the posttranslational modification of proteins produced in the rough ER while assisting in the transport of proteins through the Golgi. The human GP73 gene has been mapped within a BAC and localized to chromosome 9q21.33. GP73 levels rise in those who have been diagnosed with acute and chronic liver diseases.

## REFERENCES

1. Kladney, R.D., Bulla, G.A., Guo, L., Mason, A.L., Tollefson, A.E., Simon, D.J., Koutoubi, Z. and Fimmel, C.J. 2000. GP73, a novel Golgi-localized protein upregulated by viral infection. *Gene* 249: 53-65.
2. Kladney, R.D., Cui, X., Bulla, G.A., Brunt, E.M. and Fimmel, C.J. 2002. Expression of GP73, a resident Golgi membrane protein, in viral and nonviral liver disease. *Hepatology* 35: 1431-1440.
3. Kladney, R.D., Tollefson, A.E., Wold, W.S. and Fimmel, C.J. 2002. Upregulation of the Golgi protein GP73 by adenovirus infection requires the E1A CtBP interaction domain. *Virology* 301: 236-246.
4. Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 606804. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
5. Iftikhar, R., Kladney, R.D., Havlioglu, N., Schmitt-Gräff, A., Gusmirovic, I., Solomon, H., Luxon, B.A., Bacon, B.R. and Fimmel, C.J. 2004. Disease- and cell-specific expression of GP73 in human liver disease. *Am. J. Gastroenterol.* 99: 1087-1095.
6. Maitra, A. and Thuluvath, P.J. 2004. GP73 and liver disease: a (Golgi) complex enigma. *Am. J. Gastroenterol.* 99: 1096-1098.
7. Block, T.M., Comunale, M.A., Lowman, M., Steel, L.F., Romano, P.R., Fimmel, C., Tennant, B.C., London, W.T., Evans, A.A., Blumberg, B.S., Dwek, R.A., Mattu, T.S. and Mehta, A.S. 2005. Use of targeted glycoproteomics to identify serum glycoproteins th with liver cancer in woodchucks and humans. *Proc. Natl. Acad. Sci. USA* 102: 779-784.
8. Marrero, J.A., Romano, P.R., Nikolaeva, O., Steel, L., Mehta, A., Fimmel, C.J., Comunale, M.A., D'Amelio, A., Lok, A.S. and Block, T.M. 2005. GP73, a resident Golgi carcinoma. *J. Hepatol.* 43: 1007-1012.
9. Tsuji, A.B., Sugyo, A., Ogiu, T., Sagara, M., Kimura, T., Ishikawa, A., Sudo, H., Ohtsuki, M., Aburatani, H., Imai, T. and Harada, Y.N. 2005. Fine mapping of radiation susceptibility and gene expression analysis of LEC congenic rat lines. *Genomics* 86: 271-279.

## CHROMOSOMAL LOCATION

Genetic locus: *Golm1* (mouse) mapping to 13 B2.

## PRODUCT

GP73 (m): 293T Lysate represents a lysate of mouse GP73 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

## APPLICATIONS

GP73 (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive GP73 antibodies. Recommended use: 10-20 µl per lane.


Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

GP73 (G-2): sc-398230 is recommended as a positive control antibody for Western Blot analysis of enhanced mouse GP73 expression in GP73 transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

## RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:  
 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

## DATA



GP73 (G-2): sc-398230. Western blot analysis of GP73 expression in non-transfected: sc-117752 (A) and mouse GP73 transfected: sc-120573 (B) 293T whole cell lysates.

## STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.