



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

HA-1 (m): 293T Lysate: sc-120698

BACKGROUND

Major histocompatibility complex (MHC) molecules, which include human leukocyte antigens (HLAs), form an integral part of the immune response system. They are cell-surface receptors that bind foreign peptides and present them to cytotoxic T lymphocytes (CTLs). Minor histocompatibility antigens can form an immune response upon recognition by certain T cells when complexed with MHC molecules. HA-1 (minor histocompatibility protein HA-1), also known as HA-1, HLA-HA1 or HMHA1, is a 1,136 amino acid GTPase activator of Rho-type GTPases. Expressed in dendritic cells, epidermal Langerhans cells, hematopoietic cells, peripheral blood mononuclear cells and all leukemia and lymphoma cell lines, HA-1 is also found in various solid tissues and tumors. Highly phosphorylated, HA-1 contains one Rho-GAP domain, a single phorbol-ester/DAG-type zinc finger and is encoded by a gene located on human chromosome 19p13.3.

REFERENCES

1. van Els, C.A., D'Amato, J., Pool, J., Blokland, E., Bakker, A., van Elsen, P.J., van Rood, J.J. and Goulmy, E. 1992. Immunogenetics of human minor histocompatibility antigens: their polymorphism and immunodominance. *Immunogenetics* 35: 161-165.
2. Schreuder, G.M., Pool, J., Blokland, E., van Els, C., Bakker, A., van Rood, J.J. and Goulmy, E. 1993. A genetic analysis of human minor histocompatibility antigens demonstrates Mendelian segregation independent of HLA. *Immunogenetics* 38: 98-105.
3. Goulmy, E., Schipper, R., Pool, J., Blokland, E., Falkenburg, J.H., Vossen, J., Gratwohl, A., Vogelsang, G.B., van Houwelingen, H.C. and van Rood, J.J. 1996. Mismatches of minor histocompatibility antigens between HLA-identical donors and recipients and the development of graft-versus-host disease after bone marrow transplantation. *N. Engl. J. Med.* 334: 281-285.
4. den Haan, J.M., Meadows, L.M., Wang, W., Pool, J., Blokland, E., Bishop, T.L., Reinhardus, C., Shabanowitz, J., Offringa, R., Hunt, D.F., Engelhard, V.H. and Goulmy, E. 1998. The minor histocompatibility antigen HA-1: a diallelic gene with a single amino acid polymorphism. *Science* 279: 1054-1057.
5. Kaminski, W.E., Piehler, A. and Schmitz, G. 2000. Genomic organization of the human cholesterol-responsive ABC transporter ABCA7: tandem linkage with the minor histocompatibility antigen HA-1 gene. *Biochem. Biophys. Res. Commun.* 278: 782-789.
6. Mommaas, B., Stegehuis-Kamp, J.A., van Halteren, A.G., Kester, M., Enczmann, J., Wernet, P., Kögler, G., Mutis, T., Brand, A. and Goulmy, E. 2005. Cord blood comprises antigen-experienced T cells specific for maternal minor histocompatibility antigen HA-1. *Blood.* 105: 1823-1827.
7. Online Mendelian Inheritance in Man, OMIM™. 2009. Johns Hopkins University, Baltimore, MD. MIM Number: 601155. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

CHROMOSOMAL LOCATION

Genetic locus: Hmha1 (mouse) mapping to 10 C1.

PRODUCT

HA-1 (m): 293T Lysate represents a lysate of mouse HA-1 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

APPLICATIONS

HA-1 (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive HA-1 antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.