



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

HES1 (m2): 293T Lysate: sc-120760

BACKGROUND

The *Drosophila* Hairy and enhancer of split genes encode basic helix-loop-helix (bHLH) transcriptional repressors that function in the Notch signaling pathway and control segmentation and neural development during embryogenesis. The mammalian homolog of *Drosophila* Hairy and enhancer of split are the HES gene family members HES1-6, which also encode bHLH transcriptional repressors that regulate myogenesis and neurogenesis. The HES family members form a complex with TLE, the mammalian homolog of groucho, and this interaction is mediated by the carboxy-terminal WRPW motif of the HES proteins. The HES/TLE complex functions by directly binding to DNA instead of interfering with activator proteins. Most HES family members, including HES1 and HES5, preferentially bind to the N box (CACNAG) as opposed to the E box (CANNTG). HES1 and HES2 are expressed in a variety of adult and embryonic tissues.

REFERENCES

1. Sasai, Y., Kageyama, R., Tagawa, Y., Shigemoto, R. and Nakanishi, S. 1992. Two mammalian helix-loop-helix factors structurally related to *Drosophila* Hairy and Enhancer of split. *Genes Dev.* 6: 2620-2634.
2. Akazawa, C., Sasai, Y., Nakanishi, S. and Kageyama, R. 1992. Molecular characterization of a rat negative regulator with a basic helix-loop-helix structure predominantly expressed in the developing nervous system. *J. Biol. Chem.* 267: 21879-21885.
3. Ishibashi, M., Sasai, Y., Nakanishi, S. and Kageyama, R. 1993. Molecular characterization of HES2, a mammalian helix-loop-helix factor structurally related to *Drosophila* Hairy and Enhancer of split. *Eur. J. Biochem.* 215: 645-652.
4. Takebayashi, K., Sasai, Y., Sakai, Y., Watanabe, T., Nakanishi, S. and Kageyama, R. 1994. Structure, chromosomal locus and promoter analysis of the gene encoding the mouse helix-loop-helix factor HES1. Negative autoregulation through the multiple N box elements. *J. Biol. Chem.* 269: 5150-5156.
5. Fisher, A.L., Ohsako, S. and Caudy, M. 1996. The WRPW motif of the Hairy-related basic helix-loop-helix repressor proteins acts as a 4 amino-acid transcription repression and protein-protein interaction domain. *Mol. Cell Biol.* 16: 2670-2677.
6. Grbavec, D. and Stifani, S. 1996. Molecular interaction between TLE1 and the carboxyl-terminal domain of HES1 containing the WRPW motif. *Biochem. Biophys. Res. Commun.* 223: 701-705.
7. Lobe, C.G. 1997. Expression of the helix-loop-helix factor, HES3, during embryo development suggests a role in early midbrain-hindbrain patterning. *Mech. Dev.* 62: 227-237.
8. Bae, S., Bessho, Y., Hojo, M. and Kageyama, R. 2000. The bHLH gene HES6, an inhibitor of HES1, promotes neuronal differentiation. *Development* 127: 2933-2943.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

CHROMOSOMAL LOCATION

Genetic locus: Hes1 (mouse) mapping to 16 B2.

PRODUCT

HES1 (m2): 293T Lysate represents a lysate of mouse HES1 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

APPLICATIONS

HES1 (m2): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive HES1 antibodies. Recommended use: 10-20 µl per lane.


Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

HES1 (F-10): sc-165996 is recommended as a positive control antibody for Western Blot analysis of enhanced mouse HES1 expression in HES1 transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

DATA



HES1 (F-10): sc-165996. Western blot analysis of HES1 expression in non-transfected: sc-117752 (A) and mouse HES1 transfected: sc-120760 (B) 293T whole cell lysates.



HES1 (A-12): sc-166378. Western blot analysis of HES1 expression in non-transfected: sc-117752 (A) and mouse HES1 transfected: sc-120760 (B) 293T whole cell lysates.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.