



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# hnRNP E1 (m): 293T Lysate: sc-120856

## BACKGROUND

Heterogeneous nuclear ribonucleoproteins (hnRNPs) constitute a set of polypeptides that contribute to mRNA transcription, pre-mRNA processing, mature mRNA transport to the cytoplasm and translation. They also bind heterogeneous nuclear RNA (hnRNA), which are the transcripts produced by RNA polymerase II. There are approximately 20 known hnRNP proteins, and their complexes are the major constituents of the spliceosome. The majority of hnRNP proteins components are localized to the nucleus; however some shuttle between the nucleus and the cytoplasm, such as hnRNP E1 and E2. hnRNP E1 may function in the cytoplasm as a translational regulatory protein, while hnRNP E2 stabilizes mRNA to enhance polioviral mRNA translation. hnRNP M is involved in pre-mRNA splicing and in stress-induced transient splicing arrest.

## REFERENCES

1. Badolato, J., et al. 1995. Identification and characterisation of a novel human RNA-binding protein. *Gene* 166: 323-327.
2. Siomi, H., et al. 1995. A nuclear localization domain in the hnRNP A1 protein. *J. Cell Biol.* 129: 551-560.
3. Gattoni, R., et al. 1996. The human hnRNP M proteins: structure and relation with early heat shock-induced splicing arrest and chromosome mapping. *Nucleic Acids Res.* 24: 2535-2542.
4. Ostareck, D.H., et al. 1997. mRNA silencing in erythroid differentiation: hnRNP K and hnRNP E1 regulate 15-lipoxygenase translation from the 3' end. *Cell* 89: 597-606.
5. Kim, J.H., et al. 2000. Protein-protein interaction among hnRNPs shuttling between nucleus and cytoplasm. *J. Mol. Biol.* 298: 395-405.
6. Melcak, I., et al. 2000. Nuclear pre-mRNA compartmentalization: trafficking of released transcripts to splicing factor reservoirs. *Mol. Biol. Cell* 11: 497-510.
7. Mahe, D., et al. 2000. Spatiotemporal regulation of hnRNP M and 2H9 gene expression during mouse embryonic development. *Biochim. Biophys. Acta* 1492: 414-424.

## CHROMOSOMAL LOCATION

Genetic locus: Pcbp1 (mouse) mapping to 6 D1.

## PRODUCT

hnRNP E1 (m): 293T Lysate represents a lysate of mouse hnRNP E1 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

## STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.


## APPLICATIONS

hnRNP E1 (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive hnRNP E1 antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

hnRNP E1 (E-2): sc-137249 is recommended as a positive control antibody for Western Blot analysis of enhanced mouse hnRNP E1 expression in hnRNP E1 transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

## DATA



hnRNP E1 (E-2): sc-137249. Western blot analysis of hnRNP E1 expression in non-transfected: sc-117752 (A) and mouse hnRNP E1 transfected: sc-120856 (B) 293T whole cell lysates.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.