



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# ING4 (m): 293T Lysate: sc-121066

## BACKGROUND

Inhibitor of growth protein (ING) family of nuclear proteins, also designated ING tumor suppressor proteins, inhibit tumor progression by modulating the transcriptional outputs of signaling pathways, which in turn regulates cell proliferation. Members of this family include ING1, ING2, ING3, ING4 and INGX. ING4 localizes to the nucleus and may inhibit tumor progression by adjusting the transcriptional output of signaling pathways which regulate cell proliferation. When complexed with RELA, ING4 can suppress brain tumor angiogenesis through transcriptional repression of RELA/NFKB3 target genes. ING4 interacts with HIF prolyl hydroxylase 2 (EGLN1), which represses the activity of hypoxia inducible factor (HIF).

## REFERENCES

- Shiseki, M., Nagashima, M., Pedoux, R.M., Kitahama-Shiseki, M., Miura, K., Okamura, S., Onogi, H., Higashimoto, Y., Appella, E., Yokota, J. and Harris, C.C. 2003. p29ING4 and p28ING5 bind to p53 and p300, and enhance p53 activity. *Cancer Res.* 63: 2373-2378.
- Zhang, X., Xu, L.S., Wang, Z.Q., Wang, K.S., Li, N., Cheng, Z.H., Huang, S.Z., Wei, D.Z. and Han, Z.G. 2004. ING4 induces G<sub>2</sub>/M cell cycle arrest and enhances the chemosensitivity to DNA-damage agents in Hep G2 cells. *FEBS Lett.* 570: 7-12.
- Garkavtsev, I., Kozin, S.V., Chernova, O., Xu, L., Winkler, F., Brown, E., Barnett, G.H. and Jain, R.K. 2004. The candidate tumour suppressor protein ING4 regulates brain tumour growth and angiogenesis. *Nature* 428: 328-332.
- Kim, S., Chin, K., Gray, J.W. and Bishop, J.M. 2004. A screen for genes that suppress loss of contact inhibition of ING4 as a candidate tumor suppressor gene in human cancer. *Proc. Natl. Acad. Sci. USA* 101: 16251-16256.
- Gunduz, M., Nagatsuka, H., Demircan, K., Gunduz, E., Cengiz, B., Ouchida, M., Tsujigiwa, H., Yamachika, E., Fukushima, K., Beder, L., Hirohata, S., Ninomiya, Y., Nishizaki, K., Shimizu, K. and Nagai, N. 2005. Frequent deletion and down-regulation of ING4, a candidate tumor suppressor gene at 12p13, in head and neck squamous cell carcinomas. *Gene* 356: 109-117.
- Ozer, A., Wu, L.C. and Bruick, R.K. 2005. The candidate tumor suppressor ING4 represses activation of the hypoxia inducible factor (HIF). *Proc. Natl. Acad. Sci. USA* 102: 7481-7486.
- Ozer, A. and Bruick, R.K. 2005. Regulation of HIF by prolyl hydroxylases: recruitment of the candidate tumor suppressor protein ING4. *Cell Cycle* 4: 1153-1156.
- Zhang, X., Wang, K.S., Wang, Z.Q., Xu, L.S., Wang, Q.W., Chen, F., Wei, D.Z. and Han, Z.G. 2005. Nuclear localization signal of ING4 plays a key role in its binding to p53. *Biochem. Biophys. Res. Commun.* 331: 1032-1038.

## CHROMOSOMAL LOCATION

Genetic locus: Ing4 (mouse) mapping to 6 F2.

## PRODUCT

ING4 (m): 293T Lysate represents a lysate of mouse ING4 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

## APPLICATIONS

ING4 (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive ING4 antibodies. Recommended use: 10-20 µl per lane.


Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

ING4 (B-8): sc-515714 is recommended as a positive control antibody for Western Blot analysis of enhanced mouse ING4 expression in ING4 transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

## RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:  
 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

## DATA



ING4 (B-8): sc-515714. Western blot analysis of ING4 expression in non-transfected: sc-117752 (A) and mouse ING4 transfected: sc-121066 (B) 293T whole cell lysates.

## STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.