



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

L-FABP (m): 293T Lysate: sc-121261

BACKGROUND

Fatty acid-binding proteins, designated FABPs, are a family of homologous cytoplasmic proteins that are expressed in a highly tissue-specific manner and play an integral role in the balance between lipid and carbohydrate metabolism. FABPs mediate fatty acid (FA) and/or hydrophobic ligand uptake, transport and targeting within their respective tissues. The mechanisms underlying these actions can give rise to both passive diffusional uptake and protein-mediated transmembrane transport of FAs. FABPs are expressed in adipocytes (A-FABP), brain (B-FABP), epithelium (E-FABP, psoriasis-associated FABP, PA-FABP), striated muscle and heart (H-FABP, mammary-derived growth inhibitor or MDGI), intestine (I-FABP), liver (L-FABP), myelin (M-FABP) and testis (T-FABP). Liver-specific FABP (L-FABP) expression is modulated by developmental, hormonal, dietary and pharmacological factors, and is required for cholesterol synthesis and metabolism.

REFERENCES

1. Kaikaus, R.M., Chan, W.K., Ortiz de Montellano, P.R. and Bass, N.M. 1993. Mechanisms of regulation of liver fatty acid-binding protein. *Mol. Cell. Biochem.* 123: 93-100.
2. Veerkamp, J.H. and Maatman, R.G. 1995. Cytoplasmic fatty acid-binding proteins: their structure and genes. *Prog. Lipid Res.* 34: 17-52.
3. Hotamisligil, G.S., Johnson, R.S., Distel, R.J., Ellis, R., Papanioannou, V.E. and Spiegelman, B.M. 1996. Uncoupling of obesity from Insulin resistance through a targeted mutation in aP2, the adipocyte fatty acid binding protein. *Science* 274: 1377-1379.
4. Murphy, E.J. 1998. L-FABP and I-FABP expression increase NBD-stearate uptake and cytoplasmic diffusion in L cells. *Am. J. Physiol.* 275: 244-249.
5. Storch, J. and Thumser, A.E. 2000. The fatty acid transport function of fatty acid-binding proteins. *Biochim. Biophys. Acta* 1486: 28-44.
6. Glatz, J.F. and Storch, J. 2001. Unravelling the significance of cellular fatty acid-binding proteins. *Curr. Opin. Lipidol.* 12: 267-274.
7. Veerkamp, J.H. and Zimmerman, A.W. 2001. Fatty acid-binding proteins of nervous tissue. *J. Mol. Neurosci.* 16: 133-142.
8. Online Mendelian Inheritance in Man, OMIM™. 2001. Johns Hopkins University, Baltimore, MD. MIM Number: 134640. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>

CHROMOSOMAL LOCATION

Genetic locus: *Fabp1* (mouse) mapping to 6 C1.

PRODUCT

L-FABP (m): 293T Lysate represents a lysate of mouse L-FABP transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

APPLICATIONS

L-FABP (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive L-FABP antibodies. Recommended use: 10-20 µl per lane.


Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

L-FABP (C-4): sc-374537 is recommended as a positive control antibody for Western Blot analysis of enhanced mouse L-FABP expression in L-FABP transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:
1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

DATA



L-FABP (C-4): sc-374537. Western blot analysis of L-FABP expression in non-transfected: sc-117752 (A) and mouse L-FABP transfected: sc-121261 (B) 293T whole cell lysates.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.