



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Laminin β -2 (m): 293T Lysate: sc-121282

BACKGROUND

The laminins comprise a growing family of disulfide-linked heterotrimers consisting of three genetically distinct polypeptide chains, designated α , β and γ . A major component of the basal lamina, laminins play a crucial role in providing a scaffolding upon which tissues are assembled and which serves as a physical barrier separating specialized tissues. During embryogenesis and early development, cells migrate along basement membranes, which are required for the polarization of cells. The Laminin β -2 chain plays a role in melanoma spread, promoting tumor migration along the abluminal surface of a vessel, a phenomenon which has been termed extra-vascular migratory metastasis.

REFERENCES

1. Yurchenco, P.D. and O'Rear, J.J. 1994. Basal lamina assembly. *Curr. Opin. Cell Biol.* 6: 674-681.
2. Engvall, E. 1995. Structure and function of basement membranes. *Int. J. Dev. Biol.* 39: 781-787.
3. Aumailley, M. and Krieg, T. 1996. Laminins: a family of diverse multifunctional molecules of basement membranes. *J. Invest. Dermatol.* 106: 209-214.
4. Nomizu, M., Utani, A., Beck, K., Otaka, A., Roller, P.P. and Yamada, Y. 1996. Mechanism of Laminin chain assembly into a triple-stranded coiled-coil structure. *Biochemistry* 35: 2885-2893.
5. Ziober, B.L., Lin, C.S. and Kramer, R.H. 1996. Laminin-binding integrins in tumor progression and metastasis. *Semin. Cancer Biol.* 7: 119-128.
6. Ancsin, J.B. and Kisilevsky, R. 1996. Laminin interactions important for basement membrane assembly are promoted by zinc and implicate Laminin zinc finger-like sequences. *J. Biol. Chem.* 271: 6845-6851.
7. Durkin, M.E., Gautam, M., Loechel, F., Sanes, J.R., Merlie, J.P., Albrechtsen, R. and Wewer, U.M. 1996. Structural organization of the human and mouse Laminin β -2 chain genes, and alternative splicing at the 5' end of the human transcript. *J. Biol. Chem.* 271: 13407-13416.
8. Vogel, W., Kanz, L., Brugger, W., Berndt, A. and Kosmehl, H. 1999. Expression of Laminin β -2 chain in normal human bone marrow. *Blood* 94: 1143-1145.
9. Libby, R.T., Lavalley, C.R., Balkema, G.W., Brunken, W.J. and Hunter, D.D. 1999. Disruption of Laminin β -2 chain production causes alterations in morphology and function in the CNS. *J. Neurosci.* 19: 9399-9411.

CHROMOSOMAL LOCATION

Genetic locus: Lamb2 (mouse) mapping to 9 F2.

PRODUCT

Laminin β -2 (m): 293T Lysate represents a lysate of mouse Laminin β -2 transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

APPLICATIONS

Laminin β -2 (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive Laminin β -2 antibodies. Recommended use: 10-20 μ l per lane.


Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

Laminin β -2 (C4): sc-59980 is recommended as a positive control antibody for Western Blot analysis of enhanced mouse Laminin β -2 expression in Laminin β -2 transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:
1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

DATA



Laminin β -2 (C4): sc-59980. Western blot analysis of Laminin β -2 expression in non-transfected: sc-117752 (A) and mouse Laminin β -2 transfected: sc-121282 (B) 293T whole cell lysates.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.