



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



Matrilin-3 (m): 293T Lysate: sc-121531

BACKGROUND

The matrilin family of secreted extracellular matrix proteins is comprised of Matrilin-1 through Matrilin-4. Matrilin-1 is a homotrimer that binds to collagen and is a component of the extracellular matrix of nonarticular cartilage. It is secreted primarily by chondrocytes in a characteristic spatial, temporal and developmental stage-specific pattern during skeletogenesis. Matrilin-2 is a secreted protein involved in matrix assembly. Matrilin-3 is a secreted protein expressed solely in cartilaginous tissues. It is important in the extracellular matrix of cartilage and in the formation of extracellular filamentous networks. Matrilin-4, expressed in embryonic kidney, lung and placenta, is a secreted protein important to the extracellular matrix of cartilage.

REFERENCES

1. Deak, F., et al. 1999. The matrilins: a novel family of oligomeric extracellular matrix proteins. *Matrix Biol.* 18: 55-64.
2. Segat, D., et al. 2000. Expression of Matrilin-1, -2 and -3 in developing mouse limbs and heart. *Matrix Biol.* 19: 649-655.
3. Strusberg, I., et al. 2002. Association analysis of genotypic frequencies of Matrilin-1 gene in patients with osteoarthritis. *Clin. Exp. Rheumatol.* 20: 543-545.
4. Wiberg, C., et al. 2003. Complexes of Matrilin-1 and Biglycan or Decorin connect collagen VI microfibrils to both collagen II and aggrecan. *J. Biol. Chem.* 278: 37698-37704.
5. Ohno, S., et al. 2003. Immunohistochemical study of Matrilin-1 in arthritic articular cartilage of the mandibular condyle. *J. Oral Pathol. Med.* 32: 237-242.
6. Mann H.H., et al. 2004. Interactions between the cartilage oligomeric matrix protein and matrilins. Implications for matrix assembly and the pathogenesis of chondrodysplasias. *J. Biol. Chem.* 279: 25294-25298.
7. Karcagi, I., et al. 2004. Functional analysis of the regulatory regions of the Matrilin-1 gene in transgenic mice reveals modular arrangement of tissue-specific control elements. *Matrix Biol.* 22: 605-618.
8. Hansson, A.S., et al. 2004. Critical role of the major histocompatibility complex and IL-10 in Matrilin-1-induced relapsing polychondritis in mice. *Arthritis Res. Ther.* 6: 484-491.
9. Hansson, A.S., et al. 2004. Relapsing polychondritis, induced in mice with Matrilin-1, is an antibody- and complement-dependent disease. *Am. J. Pathol.* 164: 959-966.

CHROMOSOMAL LOCATION

Genetic locus: *Matn3* (mouse) mapping to 12 A1.1.

PRODUCT

Matrilin-3 (m): 293T Lysate represents a lysate of mouse Matrilin-3 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

Matrilin-3 (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive Matrilin-3 antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.