

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com

SANTA CRUZ BIOTECHNOLOGY, INC.

MEF-2C (m): 293T Lysate: sc-121587



BACKGROUND

The myocyte enhancer factor-2 (MEF-2) family of transcription factors associate with corepessors or co-activators to regulate development and function of T cells, neuronal cells and muscle cells. Four family members arise from alternatively spliced transcripts, termed MEF-2A, -2B, -2C and -2D. These members bind as homo- and heterodimers to the MEF-2 site in the promoter region of affected genes. Differential regulation in the expression of the four transcripts implies functional distinction for each duing embryogenesis and development. The process of differentiation from mesodermal precursor cells to myoblasts has led to the discovery of a variety of tissue-specific factors that regulate muscle gene expression. The myogenic basic helix-loop-helix proteins, including MyoD, myogenin, Myf-5, and MRF4, are one class of identified factors. A second family of DNA binding regulatory proteins is the myocyte-specific enhancer factor-2 (MEF-2) family. Each of these proteins binds to the MEF-2 target DNA sequence present in the regulatory regions of many muscle-specific genes.

REFERENCES

- Hidaka, K., et al. 1995. The MEF-2B homologue differentially expressed in mouse embryonal carcinoma cells. Biochem. Biophys. Res. Commun. 213: 555-560.
- Hobson, G.M., et al. 1995. Regional chromosomal assignments for four members of the MADS domain transcription enhancer factor 2 (MEF2) gene family to human chromosomes 15q26, 19p12, 5q14, and 1q12-q23. Genomics 29: 704-711.
- Zhao, M., et al. 1999. Regulation of the MEF2 family of transcription factors by p38. Mol. Cell. Biol. 19: 21-30.
- Han, A., et al. 2003. Sequence-specific recruitment of transcriptional co-repressor Cabin-1 by myocyte enhancer factor-2. Nature 422: 730-734.
- Pal, R., et al. 2006. Role of Smad- and Wnt-dependent pathways in embryonic cardiac development. Stem Cells Dev. 15: 29-39.
- Gude, N., et al. 2006. Akt promotes increased cardiomyocyte cycling and expansion of the cardiac progenitor cell population. Circ. Res. 99: 381-388.
- Chen, S.E., et al. 2007. TNFα regulates myogenesis and muscle regeneration by activating p38 MAPK. Am. J. Physiol. Cell Physiol. 292: C1660-1671.
- Meissner, J.D., et al. 2007. The p38α/β mitogen-activated protein kinases mediate recruitment of CREB-binding protein to preserve fast myosin heavy chain IId/X gene activity in myotubes. J. Biol. Chem. 282: 7265-7275.
- Dwivedi, Y., et al. 2007. Aberrant extracellular signal-regulated kinase (ERK) 5 signaling in hippocampus of suicide subjects. Neuropsychopharmacology 32: 2338-2350.

CHROMOSOMAL LOCATION

Genetic locus: Mef2c (mouse) mapping to 13 C3.

PRODUCT

MEF-2C (m): 293T Lysate represents a lysate of mouse MEF-2C transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

APPLICATIONS

MEF-2C (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive MEF-2C antibodies. Recommended use: 10-20 μI per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.