



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# Myosin If (m): 293T Lysate: sc-121895

## BACKGROUND

Actin is a highly conserved protein that is expressed in all eukaryotic cells. Actin filaments can form both stable and labile structures and are crucial components of microvilli and the contractile apparatus of muscle cells. Troponin facilitates interaction between Actin and Myosin by binding to  $Ca^{2+}$ . Troponin is made up of at least two subunits, which are divergent in cardiac muscle, fast skeletal muscle and slow skeletal muscle. Myosin is a hexamer of two heavy chains (MHC) and four light chains (MLC) that interacts with Actin to generate the force for diverse cellular movements, including cytokinesis, phagocytosis and muscle contraction. Myosin If (MYO1F), also designated Myosin-IE, is considered an unconventional Myosin and is expressed in the cochlea. The MYO1F gene encoding for the 1,098 amino acid protein maps to chromosome 19p13.3-13.2.

## REFERENCES

1. Bárány, M. 1967. ATPase activity of Myosin correlated with speed of muscle shortening. *J. Gen. Physiol.* 50: 197-218.
2. Billeter, R., Weber, H., Lutz, H., Howald, H., Eppenberger, HM. and Jenny, E. 1980. Myosin types in human skeletal muscle fibers. *Histochemistry* 65: 249-259.
3. Whalen, R.G., Schwartz, K., Bouveret, P., Sell, S.M. and Gros, F. 1980. Contractile protein isozymes in muscle development: identification of an embryonic form of Myosin heavy chain. *Proc. Natl. Acad. Sci. USA* 76: 5197-5201.
4. Barton, P.J. and Buckingham, M.E. 1985. The Myosin alkali light chain proteins and their genes. *Biochem. J.* 231: 249-261.
5. Warrick, H.M. and Spudich, J.A. 1988. Myosin structure and function in cell motility. *Annu. Rev. Cell Biol.* 3: 379-421.
6. Crozet, F., et al. 1997. Cloning of the genes encoding two murine and human cochlear unconventional type I Myosins. *Genomics* 40: 332-341.
7. Hodge, T. and Cope, M.J. 2000. A Myosin family tree. *J. Cell Sci.* 113: 3353-3354.
8. Krugmann, S., et al. 2002. Identification of ARAP3, a novel PI3K effector regulating both Arf and Rho GTPases, by selective capture on phosphoinositide affinity matrices. *Mol. Cell* 9: 95-108.
9. Dumont, R.A., et al. 2002. Myosin I isozymes in neonatal rodent auditory and vestibular epithelia. *J. Assoc. Res. Otolaryngol.* 3: 375-389.

## CHROMOSOMAL LOCATION

Genetic locus: Myo1f (mouse) mapping to 17 B1.

## PRODUCT

Myosin If (m): 293T Lysate represents a lysate of mouse Myosin If transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

## STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

## APPLICATIONS

Myosin If (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive Myosin If antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.