

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



NEDD4-L (m2): 293T Lysate: sc-121991



The Power to Ouestion

BACKGROUND

NEDD4-L (neural precursor cell expressed, developmentally down-regulated 4-like), also known as RSP5, NEDD4-2 or NEDL3, is a 975 amino acid protein that localizes to the cytoplasm and contains 4 WW domains, one HECT domain and one C2 domain. Expressed ubiquitously with highest expression in pancreas, prostate and kidney, NEDD4-L functions as an E3 ubiquitin-protein ligase that, characteristic of E3 ligase proteins, accepts ubiquitin (in the form of a thioester) from an E2 ubiquitin-conjugating enzyme and transfers that ubiquitin residue to substrates targeted for degradation. Through its ability to ubiquitinate and induce the proteasome-dependent degradation of proteins such as Smad2 and TGF β RII, NEDD4-L is thought to inhibit the TGF β signaling pathway, thereby regulating the signaling pathways that control cell growth and differentiation. NEDD4-L is expressed as eight isoforms due to alternative splicing events.

REFERENCES

- Chen, H., et al. 2001. NEDD4-L on human chromosome 18q21 has multiple forms of transcripts and is a homologue of the mouse NEDD4-2 gene. Eur. J. Hum. Genet. 9: 922-930.
- Harvey, K.F., et al. 2002. N4WBP5, a potential target for ubiquitination by the NEDD4 family of proteins, is a novel Golgi-associated protein. J. Biol. Chem. 277: 9307-9317.
- Qi, H., et al. 2003. Androgens differentially regulate the expression of NEDD4-L transcripts in LNCaP human prostate cancer cells. Mol. Cell. Endocrinol. 210: 51-62.
- van Bemmelen, M.X., et al. 2004. Cardiac voltage-gated sodium channel Nav1.5 is regulated by NEDD4-2 mediated ubiquitination. Circ. Res. 95: 284-291.
- Hryciw, D.H., et al. 2004. Nedd4-2 functionally interacts with CIC-5: involvement in constitutive albumin endocytosis in proximal tubule cells. J. Biol. Chem. 279: 54996-55007.
- Ichimura, T., et al. 2005. 14-3-3 proteins modulate the expression of epithelial Na+ channels by phosphorylation-dependent interaction with Nedd4-2 ubiquitin ligase. J. Biol. Chem. 280: 13187-13194.
- 7. Zhou, R., et al. 2007. Nedd4-2 catalyzes ubiquitination and degradation of cell surface ENaC. J. Biol. Chem. 282: 20207-20212.
- 8. Dahlberg, J., et al. 2007. Polymorphism in NEDD4L is associated with increased salt sensitivity, reduced levels of P-renin and increased levels of Nt-proANP. PLoS ONE 2: e432.
- Araki, N., et al. 2008. Expression, transcription, and possible antagonistic interaction of the human Nedd4L gene variant: implications for essential hypertension. Hypertension 51: 773-777.

CHROMOSOMAL LOCATION

Genetic locus: Nedd4l (mouse) mapping to 18 E1.

PRODUCT

NEDD4-L (m2): 293T Lysate represents a lysate of mouse NEDD4-L transfected 293T cells and is provided as 100 μg protein in 200 μl SDS-PAGE buffer.

APPLICATIONS

NEDD4-L (m2): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive NEDD4-L antibodies. Recommended use: 10-20 μ l per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

Santa Cruz Biotechnology, Inc. 1.800.457.3801 831.457.3800 fax 831.457.3801 Europe +00800 4573 8000 49 6221 4503 0 www.scbt.com