

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



NF-1 (m2): 293T Lysate: sc-122030



The Power to Question

BACKGROUND

NF-1, also designated CTF, consists of a family of CCAAT box binding proteins that stimulate DNA replication and activate transcription. Analysis of human NF-1 messenger RNA has revealed two forms of the NF-1 protein arising from an alternate splicing of a single NF-1 gene. NF-1 binds its consensus DNA element as a homodimer via an amino-terminal DNA binding domain, and activates transcription through a putatively novel, proline-rich, carboxy-terminal transactivation domain. The NF-1 protein has been shown to recognize and bind the adenovirus type 2 promoter and activate transcription of herpes simplex virus thymidine kinase genes. The NF-1 consensus element has been found in the upstream promoter region of myriad eukaryotic genes, including that of Ha-Ras, alpha-globin, HSP 70, GRP 78, Histone H1, Myelin basic protein and in the *Xenopus laevis* vitellogenin gene promoter.

REFERENCES

- 1. Jones, K.A., et al. 1987. A cellular DNA-binding protein that activates eukaryotic transcription and DNA replication. Cell 48: 79-89.
- Morgan, W.D., et al. 1987. Two transcriptional activators, CCAAT-boxbinding transcription factor and heat shock transcription factor, interact with a human HSP 70 gene promoter. Mol. Cell. Biol. 7: 1129-1138.
- 3. Santoro, C., et al. 1988. A family of CCAAT-box-binding proteins active in transcription and DNA replication: cloning and expression of multiple cDNAs. Nature 334: 218-224.
- Mermod, N., et al. 1989. The proline-rich transcriptional activator of CTF/ NF-1 is distinct from the replication and DNA binding domain. Cell 58: 741-753.
- Inoue, T., et al. 1990. Isolation of complementary DNAs encoding a cerebellum-enriched nuclear factor I family that activates transcription from the mouse Myelin basic protein promoter. J. Biol. Chem. 265: 19065-19070.
- Wooden, S.K., et al. 1991. Transactivation of the GRP 78 promoter by malfolded proteins, glycosylation block, and calcium ionophore is mediated through a proximal region containing a CCAAT motif which interacts with CTF/NF-I. Mol. Cell. Biol. 11: 5612-5623.
- Dusserre, Y., et al. 1992. Purified cofactors and Histone H1 mediate transcriptional regulation by CTF/NF-1. Mol. Cell. Biol. 12: 5228-5237.
- Cardinaux, J.R., et al. 1994. Complex organization of CTF/NF-I, C/EBP, and HNF3 binding sites within the promoter of the liver-specific vitellogenin gene. J. Biol. Chem. 269: 32947-32956.
- 9. Degterev, A. et al. 1999. The role of NF-1 factors in regulation of elastin gene transcription. Matrix Biol. 18(3):295-307. PMID: 10429948

PRODUCT

NF-1 (m): 293T Lysate represents a lysate of mouse NF-1 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

APPLICATIONS

NF-1 (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive NF-1 antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

Santa Cruz Biotechnology, Inc. 1.800.457.3801 831.457.3801 fax 831.457.3801 Europe +00800 4573 8000 49 6221 4503 0 www.scbt.com