



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Nup93 (m): 293T Lysate: sc-122177

BACKGROUND

The nuclear pore complex (NPC) mediates bidirectional macromolecular traffic between the nucleus and cytoplasm in eukaryotic cells and is comprised of more than 100 different subunits. Many of the subunits belong to a family called nucleoporins (Nups), which are characterized by the presence of O-linked-N-acetylglucosamine moieties and a distinctive pentapeptide repeat (XFXFG). Nup93 (nucleoporin 93) is the most abundant nucleoporin found per NPC, contributing over 10% of the mass. It localizes to the nuclear side of the NPC, predominantly in the basket terminal ring area, and exists in a complex with Nup188, Nup53 and Nup205. This complex is crucial for NPC stability and proper assembly. Nup93 interacts directly with the Nup62 complex located at the center of the NPC and thus tethers the two subcomplexes. Nup93 is composed of a coiled-coil domain at its N-terminus and a C-terminal helical domain. Its proper function is essential for cell viability and normal NPC function.

REFERENCES

- Grandi, P., Dang, T., Pané, N., Shevchenko, A., Mann, M., Forbes, D. and Hurt, E. 1997. Nup93, a vertebrate homologue of yeast Nic96p, forms a complex with a novel 205-kDa protein and is required for correct nuclear pore assembly. *Mol. Biol. Cell* 8: 2017-2038.
- Kosova, B., Panté, N., Rollenhagen, C. and Hurt, E. 1999. Nup192p is a conserved nucleoporin with a preferential location at the inner site of the nuclear membrane. *J. Biol. Chem.* 274: 22646-22651.
- Hase, M.E. and Cordes, V.C. 2003. Direct interaction with Nup153 mediates binding of TPR to the periphery of the nuclear pore complex. *Mol. Biol. Cell* 14: 1923-1940.
- Krull, S., Thyberg, J., Björkroth, B., Rackwitz, H.R. and Cordes, V.C. 2004. Nucleoporins as components of the nuclear pore complex core structure and TPR as the architectural element of the nuclear basket. *Mol. Biol. Cell* 15: 4261-4277.
- Galy, V., Mattaj, I.W. and Askjaer, P. 2003. *Caenorhabditis elegans* nucleoporins Nup93 and Nup205 determine the limit of nuclear pore complex size exclusion *in vivo*. *Mol. Biol. Cell* 14: 5104-5115.
- Hawryluk-Gara, L.A., Shibuya, E.K. and Wozniak, R.W. 2005. Vertebrate Nup53 interacts with the nuclear lamina and is required for the assembly of a Nup93-containing complex. *Mol. Biol. Cell* 16: 2382-2394.
- Ryan, C.M., Harries, J.C., Kindle, K.B., Collins, H.M. and Heery, D.M. 2006. Functional interaction of CREB binding protein (CBP) with nuclear transport proteins and modulation by HDAC inhibitors. *Cell Cycle* 5: 2146-2152.
- Shang, X., He, Y., Zhang, L., Chen, B., He, C.J., Cheng, H.H. and Zhou, R.J. 2006. Molecular cloning of the rice field eel Nup93 with predominant expression in gonad and kidney. *Yi Chuan Xue Bao* 33: 41-48.
- Judy, S. and Schwartz, T.U. 2007. Crystal structure of nucleoporin Nic96 reveals a novel, intricate helical domain architecture. *J. Biol. Chem.* 282: 34904-34912.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

CHROMOSOMAL LOCATION

Genetic locus: Nup93 (mouse) mapping to 8 C5.

PRODUCT

Nup93 (m): 293T Lysate represents a lysate of mouse Nup93 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.


APPLICATIONS

Nup93 (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive Nup93 antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

Nup93 (3332C2a): sc-81343 is recommended as a positive control antibody for Western Blot analysis of enhanced mouse Nup93 expression in Nup93 transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

DATA



Nup93 (3332C2a): sc-81343. Western blot analysis of Nup93 expression in non-transfected: sc-117752 (A) and mouse Nup93 transfected: sc-122177 (B) 293T whole cell lysates.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.