



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

PACSIN2 (m): 293T Lysate: sc-122349

BACKGROUND

PACSINs are members of a family of cytoplasmic adapter proteins, which share a conserved C-terminal protein binding SH3 domain and a CDC15-NT domain. PACSIN1-related proteins include syndapin 1 (the rat homolog of PACSIN 1), FAP52, EM13, and PSTPIP, all of which seem to be involved in signaling pathways associated with cytoskeletal organization. PACSIN1 expression is restricted to terminally differentiated neural tissue, whereas PACSIN2 is widely expressed. PACSIN2 shows vesicle-like distribution and may be involved in regulating endocytotic processes.

REFERENCES

1. Frosch, P.M., Geier, C., Kaup, F.J., Muller, A. and Frosch, M. 1993. Molecular cloning of an echinococcal microtrichal antigen immunoreactive in Echinococcus multilocularis disease. *Mol. Biochem. Parasitol.* 58: 301-310.
2. Merilainen, J., Lehto, V.P. and Wasenius, V.M. 1997. FAP52, a novel, SH3 domain-containing focal adhesion protein. *J. Biol. Chem.* 272: 23278-23284.
3. Wu, Y., Spencer, S.D. and Lasky, L.A. 1998. Tyrosine phosphorylation regulates the SH3-mediated binding of the Wiskott-Aldrich syndrome protein to PSTPIP, a cytoskeletal-associated protein. *J. Biol. Chem.* 273: 5765-5770.
4. Plomann, M., Lange, R., Vopper, G., Cremer, H., Heinlein, U.A., Scheff, S., Baldwin, S.A., Leitges, M., Cramer, M., Paulsson, M. and Barthels, D. 1998. PACSIN, a brain protein that is upregulated upon differentiation into neuronal cells. *Eur. J. Biochem.* 256: 201-211.
5. Ritter, B., Modregger, J., Paulsson, M. and Plomann, M. 1999. PACSIN 2, a novel member of the PACSIN family of cytoplasmic adapter proteins. *FEBS Lett.* 454: 356-362.
6. Qualmann, B., Roos, J., DiGregorio, P.J. and Kelly, R.B. 1999. Syndapin I, a synaptic dynamin-binding protein that associates with the neural Wiskott-Aldrich syndrome protein. *Mol. Biol. Cell* 10: 501-513.

CHROMOSOMAL LOCATION

Genetic locus: Paccin2 (mouse) mapping to 15 E1.

PRODUCT

PACSIN2 (m): 293T Lysate represents a lysate of mouse PACSIN2 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

APPLICATIONS

PACSIN2 (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive PACSIN2 antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.