



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



PAFAH1B2 (m): 293T Lysate: sc-122352

BACKGROUND

PAFAH1B2 (platelet-activating factor acetylhydrolase IB subunit β), also known as PAFAHB, is a 229 amino acid cytoplasmic protein that belongs to the GDSL lipolytic enzyme family and the platelet-activating factor acetylhydrolase IB β/γ subunits subfamily. A ubiquitously expressed catalytic subunit of the cytosolic PAFAH1B heterotrimeric complex, PAFAH1B2 inactivates PAF by removing the acetyl group at the sn-2 position. Along with the β subunit, PAFAH1B is made up of α and γ subunits. The gene that encodes PAFAH1B2 consists of approximately 32,628 bases and maps to human chromosome 11q23.3. Chromosome 11 houses over 1,400 genes and comprises nearly 4% of the human genome. Jervell and Lange-Nielsen syndrome, Jacobsen syndrome, Niemann-Pick disease, hereditary angioedema and Smith-Lemli-Opitz syndrome are associated with defects in genes that map to chromosome 11.

REFERENCES

- Albrecht, U., Abu-Issa, R., Rätz, B., Hattori, M., Aoki, J., Arai, H., Inoue, K. and Eichele, G. 1996. Platelet-activating factor acetylhydrolase expression and activity suggest a link between neuronal migration and platelet-activating factor. *Dev. Biol.* 180: 579-593.
- Adachi, H., Tsujimoto, M., Hattori, M., Arai, H. and Inoue, K. 1997. Differential tissue distribution of the β - and γ -subunits of human cytosolic platelet-activating factor acetylhydrolase (isoform I). *Biochem. Biophys. Res. Commun.* 233: 10-13.
- Moro, F., Arrigo, G., Fogli, A., Bernard, L. and Carrozzo, R. 1998. The β and γ subunits of the human platelet-activating factor acetyl hydrolase isoform Ib (PAFAH1B2 and PAFAH1B3) map to chromosome 11q23 and 19q13.1, respectively. *Genomics* 51: 157-159.
- Online Mendelian Inheritance in Man, OMIM™. 1998. Johns Hopkins University, Baltimore, MD. MIM Number: 602508. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
- Sweeney, K.J., Clark, G.D., Prokscha, A., Dobyns, W.B. and Eichele, G. 2000. Lissencephaly associated mutations suggest a requirement for the PAFAH1B heterotrimeric complex in brain development. *Mech. Dev.* 92: 263-271.
- Sheffield, P.J., McMullen, T.W., Li, J., Ho, Y.S., Garrard, S.M., Derewenda, U. and Derewenda, Z.S. 2001. Preparation and crystal structure of the recombinant $\alpha 1/\alpha 2$ catalytic heterodimer of bovine brain platelet-activating factor acetylhydrolase Ib. *Protein Eng.* 14: 513-519.
- Hasstedt, S.J., Scott, B.T., Rosendaal, F.R., Callas, P.W., Vossen, C.Y., Long, G.L. and Bovill, E.G. 2007. Exclusion of the $\alpha 2$ subunit of platelet-activating factor acetylhydrolase 1 β (PAFAH1B2) as a prothrombotic gene in a protein C-deficient kindred and population-based case-control sample. *Thromb. Haemost.* 98: 587-592.
- Scott, B.T., Olson, N., Long, G.L. and Bovill, E.G. 2008. Novel isoforms of intracellular platelet activating factor acetylhydrolase (PAFAH1b2) in human testis; encoded by alternatively spliced mRNAs. *Prostaglandins Other Lipid Mediat.* 85: 69-80.

CHROMOSOMAL LOCATION

Genetic locus: Pafah1b2 (mouse) mapping to 9 A5.2.

PRODUCT

PAFAH1B2 (m): 293T Lysate represents a lysate of mouse PAFAH1B2 transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

APPLICATIONS

PAFAH1B2 (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive PAFAH1B2 antibodies. Recommended use: 10-20 μ l per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.


PAFAH1B2 (E-9): sc-393217 is recommended as a positive control antibody for Western Blot analysis of enhanced mouse PAFAH1B2 expression in PAFAH1B2 transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:

1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

DATA



PAFAH1B2 (E-9): sc-393217. Western blot analysis of PAFAH1B2 expression in non-transfected: sc-117752 (**A**) and mouse PAFAH1B2 transfected: sc-122352 (**B**) 293T whole cell lysates.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.