



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Pitx1 (m): 293T Lysate: sc-122598

BACKGROUND

Pitx1 and Pitx2 are highly homologous, bicoid-related transcription factors. Pitx1 is a bicoid-related homeodomain factor that exhibits preferential expression in the hindlimb, as well as expression in the developing anterior pituitary gland and first branchial arch. Deletion of the Pitx1 locus results in decreased distal expression of the hindlimb-specific marker, the T-box factor (TBX4). Pitx1 may modulate morphogenesis, growth and patterning of a specific hindlimb region, and serves as a component of the variables that influence morphological and growth distinctions in forelimb and hindlimb identity. Pitx2 was initially identified as the gene responsible for human Rieger syndrome, an autosomal dominant condition that causes developmental abnormalities. Pitx2 is a transcription factor that regulates cardiac positioning and pituitary and tooth morphogenesis. Pitx2 also regulates lung symmetry by encoding "leftness" of the lung. Pitx2 is asymmetrically expressed in the left lateral-plate mesoderm, and mutant mice with laterality defects show altered patterns of Pitx2 expression that correlate with changes in the visceral symmetry. The genes which encode Pitx1 and Pitx2 map to human chromosomes 5q31.1 and 4q25, respectively.

REFERENCES

1. Crawford, M.J., Lanctot, C., Tremblay, J.J., Jenkins, N., Gilbert, D., Copeland, N., Beatty, B. and Drouin, J. 1997. Human and murine PTX1/Ptx1 gene maps to the region for Treacher Collins syndrome. *Mamm. Genome* 8: 841-845.
2. Gage, P.J. and Camper, S.A. 1997. Pituitary homeobox 2, a novel member of the bicoid-related family of homeobox genes, is a potential regulator of anterior structure formation. *Hum. Mol. Genet.* 6: 457-464.
3. Lin, C., Kioussi, C., O'Connell, S., Briata, P., Szeto, D., Liu, F., Izpisua-Belmonte, J. and Rosenfeld, M. 1999. Pitx2 regulates lung asymmetry, cardiac positioning and pituitary and tooth morphogenesis. *Nature* 401: 279-282.
4. Kioussi, C., Carriere, C. and Rosenfeld, M. 1999. A model for the development of the hypothalamic-pituitary axis: transcribing the hypophysis. *Mech. Dev.* 81: 23-35.
5. Hollemann, T. and Pieler, T. 1999. XPitx-1: a homeobox gene expressed during pituitary and cement gland formation of *Xenopus* embryos. *Mech. Dev.* 88: 249-252.
6. Szeto, D., Rodriguez-Esteban, C., Ryan, A., O'Connell, S., Liu, F., Kioussi, C., Gleiberman, A., Izpisua-Belmonte, J. and Rosenfeld, M. 1999. Role of the Bicoid-related homeodomain factor Pitx1 in specifying hindlimb morphogenesis and pituitary development. *Genes Dev.* 13: 484-494.
7. Graham, A. and McGonnell, I. 1999. Limb development: Farewell to arms. *Curr Biol.* 9: 368-370.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

CHROMOSOMAL LOCATION

Genetic locus: Pitx1 (mouse) mapping to 13 B1.

PRODUCT

Pitx1 (m): 293T Lysate represents a lysate of mouse Pitx1 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

APPLICATIONS

Pitx1 (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive Pitx1 antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

RESEARCH USE

For research use only, not for use in diagnostic procedures.