



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Rap 1B (m3): 293T Lysate: sc-122967

BACKGROUND

Ras oncogenes encode GTP-binding proteins that are capable of transforming immortalized cells in culture. Two Ras-related human genes, designated Rap 1A and Rap 1B, encode 95% homologous proteins (namely Rap 1A and Rap 1B) that share a similar C-terminal Cys-Ali-Ali-Xaa sequence with Ras proteins and are ubiquitously expressed in mammalian tissues. The putative "effector" domain of Ras proteins, whose integrity is required for cell transformation as well as interaction with the putative effector protein GAP, is conserved in both Rap 1 proteins. Rap 1A is thought to interfere with Ras effector function by binding to Ras GAP in a GTP-dependent manner without affecting Rap 1A GTPase activity. Rap 2, another Ras-related protein, shares 60% identity with Rap 1A and exhibits a carboxy terminal CAAX motif and two upstream cysteines similar to those of the H-Ras, K-Ras and N-Ras proteins. In contrast with Rap 1A and Rap 1B, overexpression of Rap 2 does not interfere with the Ras signaling pathway.

REFERENCES

1. Pizon, V., Chardin, P., Lerosey, I., Olofsson, B. and Tavitian, A. 1988. Human cDNAs Rap 1 and Rap 2 homologous to the *Drosophila* gene Dras3 encode proteins closely related to Ras in the "effector" region. *Oncogene* 3: 201-204.
2. Pizon, V., Lerosey, I., Chardin, P. and Tavitian, A. 1988. Nucleotide sequence of a human cDNA encoding a Ras-related protein (Rap 1B). *Nucleic Acids Res.* 16: 7719.
3. Culine, S., Olofsson, B., Gosselin, S., Honore, N. and Tavitian, A. 1989. Expression of the Ras-related rap genes in human tumors. *Int. J. Cancer* 44: 990-994.
4. Kim, S., Mozoguchi, A., Kikuchi, A. and Takai, Y. 1990. Tissue and subcellular distributions of the SMG21/Rap 1/Krev-1 proteins which are partly distinct from those of c-Ras p21s. *Mol. Cell. Biol.* 10: 2645-2652.
5. Frech, M., John, J., Pizon, V., Chardin, P., Tavitian, A., Clark, R., McCormick, F. and Wittinghofer, A. 1990. Inhibition of GTPase activating protein stimulation of Ras p21 GTPase by the Krev-1 gene product. *Science* 249: 169-171.
6. Hata, Y., Kikuchi, A., Sasaki, T., Schaber, M.D., Gibbs, J.B. and Takai, Y. 1990. Inhibition of the Ras p21 GTPase-activating protein-stimulated GTPase activity of c-Ha-Ras p21 by SMG p21 having the same putative effector domain as Ras p21s. *J. Biol. Chem.* 265: 7104-7107.
7. Rubinfeld, B., Munemitsu, S., Clark, R., Conroy, L., Watt, K., Crosier, W.J., McCormick, F. and Polakis, P. 1991. Molecular cloning of a GTPase activating protein specific for the Krev-1 protein p21Rap 1. *Cell* 65: 1033-1042.
8. Béranger, F., Goud, B., Tavitian, A. and de Gunzburg, J. 1991. Association of the Ras-antagonistic Rap 1/Krev-1 proteins with Golgi complex. *Proc. Natl. Acad. Sci. USA* 88: 1606-1610.

CHROMOSOMAL LOCATION

Genetic locus: Rap1b (mouse) mapping to 10 D2.

PRODUCT

Rap 1B (m3): 293T Lysate represents a lysate of mouse Rap 1B transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

APPLICATIONS

Rap 1B (m3): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive Rap 1B antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.