



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



RAR γ (m2): 293T Lysate: sc-122976

BACKGROUND

Retinoids are metabolites of vitamin A (retinol) that are important signaling molecules during vertebrate development and tissue differentiation. Retinoic acid receptors (RARs) and retinoid X receptors (RXRs) are nuclear transcription factors that modulate the effects of retinoids (RA) on gene expression. Most retinoid forms (including all *trans* RA, 9-*cis* RA, 4oxo RA and 3,4 dihydro RA) activate RAR family members, whereas RXR family members are activated by 9-*cis*-RA only. RA binds RARs, inducing a change in receptor configuration that allows DNA binding and increased gene transcription from specific genes to occur. RAR family members, which include RAR α , RAR β and RAR γ , belong to the same class of nuclear transcription factors as thyroid hormone receptors, vitamin D3 receptor and ecdysone receptor. Retinoid receptor expression is tissue specific; the skin expresses RAR γ and RXR α . The expression of RAR γ and RXR β is somewhat decreased in lung cancers. The human RAR γ gene maps to chromosome 12q13.13.

REFERENCES

- Koelle, M.R., Talbot, W.S., Segraves, W.A., Bender, M.T., Cherbas, P. and Hogness, D.S. 1991. The *Drosophila* EcR gene encodes an ecdysone receptor, a new member of the steroid receptor superfamily. *Cell* 67: 59-77.
- Mattei, M.G., Riviere, M., Krust, A., Ingvarsson, S., Vennstrom, B., Islam, M.Q., Levan, G., Kautner, P., Zelent, A. and Chambon, P., et al. 1991. Chromosomal assignment of retinoic acid receptor (RAR) genes in the human, mouse, and rat genomes. *Genomics* 10: 1061-1069.
- Rees, J. 1992. The molecular biology of retinoic acid receptors: orphan from good family seeks home. *J. Dermatol.* 126: 97-104.
- Bhat, M.K., Ashizawa, K. and Cheng, S.Y. 1994. Phosphorylation enhances the target gene sequence-dependent dimerization of thyroid hormone receptor with retinoid X receptor. *Proc. Natl. Acad. Sci. USA* 91: 7927-7931.
- Mangelsdorf, D.J., Umesono, K. and Evans, R.M. 1994. The Retinoids. In Sporn, M.B., Roberts, A.B. and Goodman, D.S., eds., *Biology, Chemistry, and Medicine*, 2nd Edition. New York: Raven Press, Ltd., 314-349.
- Lotan, R. 1997. Roles of retinoids and their nuclear receptors in the development and prevention of upper aerodigestive tract cancers. *Environ. Health Perspect.* 105: 985-988.
- Mark, M., Kastner, P., Ghyselinck, N.B., Krezel, W., Dupe, V. and Chambon, P. 1997. Genetic control of the development by retinoic acid. *C.R. Seances Soc. Biol. Fil.* 191: 77-90.
- Orfanos, C.E., Zouboulis, C.C., Almond-Roesler, B. and Geilen, C.C. 1997. Current use and future potential role of retinoids in dermatology. *Drugs* 53: 358-388.
- Lotan, R. 1997. Retinoids and chemoprevention of aerodigestive tract cancers. *Cancer Metastasis Rev.* 16: 349-356.

CHROMOSOMAL LOCATION

Genetic locus: Rarg (mouse) mapping to 15 F3.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PRODUCT

RAR γ (m2): 293T Lysate represents a lysate of mouse RAR γ transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

APPLICATIONS

RAR γ (m2): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive RAR γ antibodies. Recommended use: 10-20 μ l per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.


RAR γ (G-1): sc-7387 is recommended as a positive control antibody for Western Blot analysis of enhanced mouse RAR γ expression in RAR γ transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:

1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

DATA



RAR γ (G-1): sc-7387. Western blot analysis of RAR γ expression in non-transfected: sc-117752 (**A**) and mouse RAR γ transfected: sc-122976 (**B**) 293T whole cell lysates.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.