



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Rho B (m2): 293T Lysate: sc-123118

BACKGROUND

The Ras p21 family of guanine nucleotide proteins has been widely studied in view of its apparent role in signal transduction pathways and high frequency of mutations in human malignancies. It is now clear, however, that the Ras proteins (H-, K- and N-Ras p21) are members of a much larger superfamily of related proteins. Six members of this family, Rap 1A, Rap 1B, Rap 2, R-Ras, Ral A and Ral B, exhibit approximately 50% amino acid homology to Ras. The five mammalian Rho proteins (Rho A, B, C, 7 and 8) are approximately 30% homologous to Ras and are expressed in a wide range of cell types. Both Ras p21 and Rho p21, as well as other members of the Ras superfamily, contain a carboxy-terminal CAAX sequence (C, cysteine; A, aliphatic amino acid; X, any amino acid) which in the case of Ras has been shown to be essential for correct localization and function.

REFERENCES

1. Liu, A.X., et al. 2001. Rho B is dispensable for mouse development, but it modifies susceptibility to tumor formation as well as cell adhesion and growth factor signaling in transformed cells. *Mol. Cell Biol.* 21: 6906-6912.
2. Liu, A., et al. 2001. Rho B is required to mediate apoptosis in neoplastically transformed cells after DNA damage. *Proc. Natl. Acad. Sci. USA* 98: 6192-6197.
3. Ader, I., et al. 2002. Rho B controls the 24 kDa FGF-2-induced radioresistance in HeLa cells by preventing post-mitotic cell death. *Oncogene* 21: 5998-6006.
4. Sandilands, E., et al. 2004. RhoB and Actin polymerization coordinate Src activation with endosome-mediated delivery to the membrane. *Dev. Cell* 7: 855-869.
5. Wheeler, A.P., et al. 2004. Why three Rho proteins? Rho A, Rho B, Rho C, and cell motility. *Exp. Cell Res.* 301: 43-49.
6. Jiang, K., et al. 2004. Akt mediates Ras downregulation of Rho B, a suppressor of transformation, invasion, and metastasis. *Mol. Cell Biol.* 24: 5565-5576.
7. Jiang, K., et al. 2004. EGFR, ErbB-2 and Ras but not Src suppress Rho B expression while ectopic expression of Rho B antagonizes oncogene-mediated transformation. *Oncogene* 23: 1136-1145.
8. Chauhan, S., et al. 2004. Androgen control of cell proliferation and cytoskeletal reorganization in human fibrosarcoma cells: role of Rho B signaling. *J. Biol. Chem.* 279: 937-944.

CHROMOSOMAL LOCATION

Genetic locus: Rhob (mouse) mapping to 12 A1.1.

PRODUCT

Rho B (m2): 293T Lysate represents a lysate of mouse Rho B transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

APPLICATIONS

Rho B (m2): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive Rho B antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.