



**SZABO  
SCANDIC**

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](http://linkedin.com/company/szaboscandic)



# RNase H1 (m): 293T Lysate: sc-123223

## BACKGROUND

The human RNase H1 enzyme is a cytoplasmic endonuclease that degrades the RNA of RNA-DNA hybrids resulting in 5'-phosphomonoester products. Human RNase H1 cleaves RNA exclusively in an RNA/DNA duplex; neither double-strand DNA nor double-strand RNA is a viable substrate. Mn<sup>2+</sup> and N-ethylmaleimide can inhibit Mg<sup>2+</sup> dependent RNase H1 activity. The RNase H1 gene is present at similar levels in all human cells and tissues, indicating that RNase H1 may be a housekeeping protein. The human RNase H1 gene maps to chromosome 2p25.3 with pseudogenes present on chromosome 17p11.2 and chromosome 1q.

## REFERENCES

- Wu, H., Lima, W. and Crooke, S. 1998. Molecular cloning and expression of cDNA for human RNase H. *Antisense Nucleic Acid Drug Dev.* 8: 53-61.
- Cerritelli, S. and Crouch, R. 1998. Cloning, expression, and mapping of ribonucleases H of human and mouse related to bacterial RNase H1. *Genomics* 53: 300-307.
- ten Asbroek, A., van Groenigen, M., Jakobs, M., Koevoets, C., Janssen, B. and Baas, F. 2002. Ribonuclease H1 maps to chromosome 2 and has at least three pseudogene loci in the human genome. *Genomics* 79: 818-23.
- Lima, W.F., Wu, H., Nichols, J.G., Manalili, S.M., Drader, J.J., Hofstadler, S.A. and Crooke, S.T. 2003. Human RNase H1 activity is regulated by a unique redox switch formed between adjacent cysteines. *J. Biol. Chem.* 278: 14906-14912.
- Lima, W.F., Wu, H., Nichols, J.G., Prakash, T.P., Ravikumar, V. and Crooke, S.T. 2003. Human RNase H1 uses one tryptophan and two lysines to position the enzyme at the 3'-DNA/5'-RNA terminus of the heteroduplex substrate. *J. Biol. Chem.* 278: 49860-49867.
- Wu, H., Lima, W.F., Zhang, H., Fan, A., Sun, H. and Crooke, S.T. 2004. Determination of the role of the human RNase H1 in the pharmacology of DNA-like antisense drugs. *J. Biol. Chem.* 279: 17181-17189.
- Lima, W.F., Nichols, J.G., Wu, H., Prakash, T.P., Migawa, M.T., Wyrzykiewicz, T.K., Bhat, B. and Crooke, S.T. 2004. Structural requirements at the catalytic site of the heteroduplex substrate for human RNase H1 catalysis. *J. Biol. Chem.* 279: 36317-36326.
- SWISS-PROT/TrEMBL (O60930). World Wide Web URL:  
<http://www.expasy.ch/sprot/sprot-top.html>

## CHROMOSOMAL LOCATION

Genetic locus: Rnaseh1 (mouse) mapping to 12 A2.

## PRODUCT

RNase H1 (m): 293T Lysate represents a lysate of mouse RNase H1 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

## STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

## APPLICATIONS

RNase H1 (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive RNase H1 antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.


RNase H1 (H-4): sc-376326 is recommended as a positive control antibody for Western Blot analysis of enhanced mouse RNase H1 expression in RNase H1 transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

## RECOMMENDED SUPPORT REAGENTS


To ensure optimal results, the following support reagents are recommended:

1) Western Blotting: use m-IgG<sub>X</sub> BP-HRP: sc-516102 or m-IgG<sub>X</sub> BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

## DATA



RNase H1 (H-4): sc-376326. Western blot analysis of RNase H1 expression in non-transfected: sc-117752 (**A**) and mouse RNase H1 transfected: sc-123223 (**B**) 293T whole cell lysates.



RNase H1 (C-8): sc-365057. Western blot analysis of RNase H1 expression in non-transfected: sc-117752 (**A**) and mouse RNase H1 transfected: sc-123223 (**B**) 293T whole cell lysates.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.