



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Rtn-1 (m): 293T Lysate: sc-123321

BACKGROUND

The Reticulon (Rtn) family consists of four members: Rtn-1 (also designated neuroendocrine-specific protein or NSP), Rtn-2 (also designated NSP-like-1), Rtn-3 (also designated NSP-like-2) and Nogo (also designated Rtn-4A). Localized on human chromosome 14q23.1, the gene encoding Rtn-1 is expressed as three isoforms: Rtn-1A (NSP-A), Rtn-1B (NSP-B) and Rtn-1C (NSP-C), which are produced by alternative splicing, are anchored to the endoplasmic reticulum in neural and neuroendocrine tissues and cells, and may be involved in neuroendocrine secretion or in membrane trafficking. In lung cancer diagnosis, Rtn-1A appears to be a reliable marker for the detection of neuroendocrine differentiation, since most of the small cell lung carcinoma (SCLC) and carcinoma tumors show expression of Rtn-1A. Rtn-1B exists as multiple forms. Expression of Rtn-1C strongly correlates with neuronal differentiation.

REFERENCES

- Senden, N.H., et al. 1994. Subcellular localization and supramolecular organization of neuroendocrine-specific protein B (NSP-B) in small cell lung cancer. *Eur. J. Cell Biol.* 65: 341-353.
- van de Velde, H.J., et al. 1994. NSP-encoded reticulons are neuroendocrine markers of a novel category in human lung cancer diagnosis. *Cancer Res.* 54: 4769-4776.
- Geisler, J.G., et al. 1998. Molecular cloning of a novel mouse gene with predominant muscle and neural expression. *Mamm. Genome* 9: 274-282.
- Roebroek, A.J., et al. 1998. cDNA cloning, genomic organization, and expression of the human RTN2 gene, a member of a gene family encoding reticulons. *Genomics* 51: 98-106.
- Hens, J., et al. 1998. Neuronal differentiation is accompanied by NSP-C expression. *Cell Tissue Res.* 292: 229-237.
- Moreira, E.F., et al. 1999. Cloning of a novel member of the reticulon gene family (RTN3): gene structure and chromosomal localization to 11q13. *Genomics* 58: 73-81.
- GrandPre, T., et al. 2000. Identification of the Nogo inhibitor of axon regeneration as a reticulon protein. *Nature* 403: 439-444.

CHROMOSOMAL LOCATION

Genetic locus: Rtn1 (mouse) mapping to 12 C3.

PRODUCT

Rtn-1 (m): 293T Lysate represents a lysate of mouse Rtn-1 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

APPLICATIONS

Rtn-1 (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive Rtn-1 antibodies. Recommended use: 10-20 µl per lane.


Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

Rtn-1 (4A66): sc-71980 is recommended as a positive control antibody for Western Blot analysis of enhanced mouse Rtn-1 expression in Rtn-1 transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:
 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

DATA



Rtn-1 (4A66): sc-71980. Western blot analysis of Rtn-1 expression in non-transfected: sc-117752 (A) and mouse Rtn-1 transfected: sc-123321 (B) 293T whole cell lysates.

RESEARCH USE

For research use only, not for use in diagnostic procedures.