



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# Sm D1 (m): 293T Lysate: sc-123635

## BACKGROUND

U1, U2, U4, U5, U6 and U7 are small nuclear ribonucleoproteins (snRNPs) that comprise the spliceosome in eukaryotes. Each UsnRNP contains common Sm proteins B/B', D1, D2, D3, E, F and G. The Sm proteins pair up as D1-D2, B/B'-D3 and E-F-G to form RNA-free hetero-oligomers in the cytoplasm. Sm proteins aid in the cytoplasmic construction of the UsnRNPs by binding to a conserved Sm site on UsnRNA and forming a stable snRNP core complex. Sm D1, D2 and D3 are present in U1, U2, U4/5 and U5 but not U7 snRNPs in human and mouse cells. U7 snRNPs contain Lsm10, an Sm D1-like protein. Autoantibodies produced in patients suffering from systemic lupus erythematosus react predominantly with Sm B/B', D1 and D3. The major linear epitope of these autoantibodies includes the C-terminal RG dipeptide repeats found in Sm D1 and D3.

## REFERENCES

1. Branlant, C., et al. 1982. U2 RNA shares a structural domain with U1, U4, and U5 RNAs. *EMBO J.* 1: 1259-1265.
2. Lehmeier, T., et al. 1990. Evidence for three distinct D proteins, which react differentially with anti-Sm autoantibodies, in the cores of the major snRNPs U1, U2, U4/U6 and U5. *Nucleic Acids Res.* 18: 6475-6484.
3. Raker, V.A., et al. 1996. The snRNP core assembly pathway: identification of stable core protein heteromeric complexes and an snRNP subcore particle *in vitro*. *EMBO J.* 15: 2256-2269.
4. Brahms, H., et al. 2000. The C-terminal RG dipeptide repeats of the spliceosomal Sm proteins D1 and D3 contain symmetrical dimethylarginines, which form a major B cell epitope for anti-Sm autoantibodies. *J. Biol. Chem.* 275: 17122-17129.
5. Pillai, R.S., et al. 2001. Purified U7 snRNPs lack the Sm proteins D1 and D2 but contain Lsm10, a new 14 kDa Sm D1-like protein. *EMBO J.* 20: 5470-5479.
6. LocusLink Report (LocusID: 6632). <http://www.ncbi.nlm.nih.gov/LocusLink/>

## CHROMOSOMAL LOCATION

Genetic locus: Snrpd1 (mouse) mapping to 18 A1.

## PRODUCT

Sm D1 (m): 293T Lysate represents a lysate of mouse Sm D1 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

## APPLICATIONS

Sm D1 (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive Sm D1 antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

## STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.