



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



Smad6 (m): 293T Lysate: sc-123643

BACKGROUND

Smad proteins, the mammalian homologs of the *Drosophila* mothers against dpp (Mad) have been implicated as downstream effectors of TGF β /BMP signaling. Smad1 (also designated Madr1 or JV4-1), Smad5 and mammalian Smad8 (also designated Smad9 or MadH6) are effectors of BMP-2 and BMP-4 function, while Smad2 (also designated Madr2 or JV18-1) and Smad3 are involved in TGF β and Activin-mediated growth modulation. Smad4 (also designated DPC4) has been shown to mediate all of the above activities through interaction with various Smad family members. Smad6 and Smad7 regulate the response to Activin/TGF β signaling by interfering with TGF β -mediated phosphorylation of other Smad family members.

REFERENCES

1. Liu, F., Hata, A., Baker, J.C., Doody, J., Carcamo, J., Harland, R.M. and Massague, J. 1996. A human Mad protein acting as a BMP-regulated transcriptional activator. *Nature* 381: 620-623.
2. Zhang, Y., Feng, X.H., Wu, R.Y. and Derynck, R. 1996. Receptor-associated Mad homologues synergize as effectors of the TGF β response. *Nature* 383: 168-172.
3. Lagna, G., Hata, A., Hemmati-Brivanlou, A. and Massague, J. 1996. Partnership between DPC4 and Smad proteins in TGF β signalling pathways. *Nature* 383: 832-836.
4. Eppert, K., Scherer, S.W., Ozcelik, H., Pirone, R., Hoodless, P., Kim, H., Tsui, L.C., Bapat, B., Gallinger, S., Andrusis, I.L., Thomsen, G.H., Wrana, J.L. and Attisano, L. 1996. Madr2 maps to 18q21 and encodes at TGF β -regulated Mad-related protein that is functionally encoded in colorectal carcinoma. *Cell* 86: 543-552.
5. Massague, J., Hata, A. and Liu, F. 1997. TGF β signalling through the Smad pathway. *Trends Cell Biol.* 7: 187-192.
6. Chen, Y., Bhushan, A. and Vale, W. 1997. Smad8 mediates the signaling of the receptor serine kinase. *Proc. Natl. Acad. Sci. USA* 94: 12938-12943.
7. Imamura, T., Takase, M., Nishihara, A., Oeda, E., Hanai, J.I., Kawabata, M. and Miyazono, K. 1997. Smad6 inhibits signalling by the TGF β superfamily. *Nature* 389: 622-626.
8. Heldin, C.H., Miyazono, K. and ten Dijke, P. 1997. TGF β signalling from cell membrane to nucleus through Smad proteins. *Nature* 390: 465-471.

CHROMOSOMAL LOCATION

Genetic locus: Smad6 (mouse) mapping to 9 C.

PRODUCT

Smad6 (m): 293T Lysate represents a lysate of mouse Smad6 transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

APPLICATIONS

Smad6 (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive Smad6 antibodies. Recommended use: 10-20 μ l per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.


Smad6 (D-4): sc-25321 is recommended as a positive control antibody for Western Blot analysis of enhanced mouse Smad6 expression in Smad6 transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:

1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

DATA



Smad6 (D-4): sc-25321. Western blot analysis of Smad6 expression in non-transfected: sc-117752 (**A**) and mouse Smad6 transfected: sc-123643 (**B**) 293T whole cell lysates.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.