



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

SMAP1 (m): 293T Lysate: sc-123645

BACKGROUND

SMAP1 (stromal membrane-associated protein 1), also known as small ArfGAP 1, is a 467 amino acid peripheral membrane protein that localizes to the cytoplasmic side of the cell membrane where it participates in clathrin-dependent endocytosis. A GTPase activating protein for ARF6, SMAP1 is widely expressed in tissues such as lymph node, spinal cord, bone marrow, adrenal gland, trachea, stomach, thyroid and embryonic hematopoietic tissues. Containing one Arf-GAP domain, SMAP1 exists as multiple isoforms as a result of alternative splicing events and is encoded by a gene that maps to human chromosome 6q13. Human chromosome 6 contains 170 million base pairs, comprises nearly 6% of the human genome and is associated with Parkinson's disease, Stickler syndrome and a susceptibility to bipolar disorder.

REFERENCES

1. Brunner, H.G., van Beersum, S.E., Warman, M.L., Olsen, B.R., Ropers, H.H. and Mariman, E.C. 1994. A Stickler syndrome gene is linked to chromosome 6 near the COL11A2 gene. *Hum. Mol. Genet.* 3: 1561-1564.
2. Sato, Y., Hong, H.N., Yanai N. and Obinata, M. 1998. Involvement of stromal membrane-associated protein (SMAP-1) in erythropoietic microenvironment. *J. Biochem.* 124: 209-216.
3. Obinata, M. and Yanai, N. 1999. Cellular and molecular regulation of an erythropoietic inductive microenvironment (EIM). *Cell Struct. Funct.* 24: 171-179.
4. Marcos, I., Borrego, S., Rodríguez de Córdoba, S., Galán, J.J. and Antiñolo, G. 2002. Cloning, characterization and chromosome mapping of the human SMAP1 gene. *Gene* 292: 167-171.
5. Cesari, R., Martin, E.S., Calin, G.A., Pentimalli, F., Bichi, R., McAdams, H., Trapasso, F., Drusco, A., Shimizu, M., Masciullo, V., D'Andrilli, G., Scambia, G., Picchio, M.C., Alder, H., Godwin A.K. and Croce, C.M. 2003. Parkin, a gene implicated in autosomal recessive juvenile parkinsonism, is a candidate tumor suppressor gene on chromosome 6q25-q27. *Proc. Natl. Acad. Sci. USA* 100: 5956-5961.
6. Barragan, I., Marcos, I., Borrego S. and Antiñolo, G. 2005. Mutation screening of three candidate genes, ELOVL5, SMAP1 and GLUL1 in autosomal recessive retinitis pigmentosa. *Int. J. Mol. Med.* 16: 1163-1167.
7. Tanabe, K., Torii, T., Natsume, W., Braesch-Andersen, S., Watanabe T., Satake, M. 2005. A novel GTPase-activating protein for ARF6 directly interacts with clathrin and regulates clathrin-dependent endocytosis. *Mol. Biol. Cell* 16: 1617-1628.

CHROMOSOMAL LOCATION

Genetic locus: Smap1 (mouse) mapping to 1 A5.

PRODUCT

SMAP1 (m): 293T Lysate represents a lysate of mouse SMAP1 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

SMAP1 (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive SMAP1 antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.